

Table of Contents


2 Poisson
11 Other Discrete RVs
19 Exercises

o8: Poisson and More



Jerry Cain
January 26, 2024

[Lecture Discussion on Ed](#)

Poisson



Algorithmic ride sharing



Probability of k requests from this area in the next 1 minute?

Suppose we know:

On average, $\lambda = 5$ requests per minute

Algorithmic ride sharing, approximately

Probability of k requests from this area in the next 1 minute?

On average, $\lambda = 5$ requests per minute

Break a minute down into 60 seconds:

0	0	1	0	1	...					0	0	0	0	1
1	2	3	4	5										60

At each second:

- Independent Bernoulli trial
- You get a request (1) or you don't (0).

Let $X = \#$ of requests in minute.

$$E[X] = \lambda = 5$$

$$X \sim \text{Bin}(n = 60, p = 5/60)$$

$$P(X = k) = \binom{60}{k} \left(\frac{5}{60}\right)^k \left(1 - \frac{5}{60}\right)^{n-k}$$




But what if there are *two* requests in the same second?

Algorithmic ride sharing, approximately

Probability of k requests from this area in the next 1 minute?

On average, $\lambda = 5$ requests per minute

Break a minute down into 60,000 **milliseconds**:



At each **millisecond**:

- Independent Bernoulli trial
- You get a request (1) or you don't (0).

Let $X = \#$ of requests in minute.

$$E[X] = \lambda = 5$$

$$X \sim \text{Bin}(n = 60000, p = \lambda/n)$$

$$P(X = k) = \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k}$$




But what if there are *two* requests
in the same **millisecond**?

Algorithmic ride sharing, approximately

Probability of k requests from this area in the next 1 minute?

On average, $\lambda = 5$ requests per minute

Break a minute down into **infinitely small** buckets:



For each time bucket:

- Independent Bernoulli trial
- You get a request (1) or you don't (0).

Let $X = \#$ of requests in minute.

$$E[X] = \lambda = 5$$

$$X \sim \text{Bin}(n, p = \lambda/n)$$

$$P(X = k) = \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k}$$

Gnarly math incoming!

Binomial in the limit

$$\lim_{n \rightarrow \infty} \left(1 - \frac{\lambda}{n}\right)^n = e^{-\lambda}$$



$$P(X = k) = \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} \quad \text{Expand} = \lim_{n \rightarrow \infty} \frac{n!}{k!(n-k)!} \frac{\lambda^k}{n^k} \frac{\left(1 - \frac{\lambda}{n}\right)^n}{\left(1 - \frac{\lambda}{n}\right)^k}$$

$$\quad \text{Rearrange} = \lim_{n \rightarrow \infty} \frac{n!}{n^k(n-k)!} \frac{\lambda^k}{k!} \frac{\left(1 - \frac{\lambda}{n}\right)^n}{\left(1 - \frac{\lambda}{n}\right)^k} \quad \text{Def natural exponent} = \lim_{n \rightarrow \infty} \frac{n!}{n^k(n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

$$\quad \text{Expand} = \lim_{n \rightarrow \infty} \frac{n(n-1) \cdots (n-k+1)}{n^k} \frac{(n-k)!}{(n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

$$\quad \text{Limit analysis + cancel} = \lim_{n \rightarrow \infty} \frac{n^k}{n^k} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{1} \quad \text{Simplify} = \frac{\lambda^k}{k!} e^{-\lambda}$$

Algorithmic ride sharing



Probability of k requests from this area in the next 1 minute?

On average, $\lambda = 5$ requests per minute

$$P(X = k) = \frac{\lambda^k}{k!} e^{-\lambda}$$

**Poisson
distribution**

Poisson Random Variable

Consider an experiment that lasts a fixed interval of time.

def A **Poisson** random variable X is the number of successes over the experiment duration, assuming **the time that each success occurs is independent** and the average # of requests over time is constant.

$$X \sim \text{Poi}(\lambda)$$

Support: $\{0, 1, 2, \dots\}$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation $E[X] = \lambda$

Variance $\text{Var}(X) = \lambda$

Examples:

- # earthquakes per year
- # server hits per second
- # of emails per day

Yes, expectation == variance for Poisson RV! More later.

Earthquakes


$$X \sim \text{Poi}(\lambda) \quad p(k) = e^{-\lambda} \frac{\lambda^k}{k!}$$
$$E[X] = \lambda$$

There are an average of 2.79 major earthquakes in the world each year, and major earthquakes occur independently.

What is the probability of 3 major earthquakes happening next year?

1. Define RVs

2. Solve





Other Discrete RVs

Grid of random variables

	Number of successes	Time until success
One trial	$\text{Ber}(p)$	One success
Several trials	$\text{Bin}(n, p)$	Several successes
Interval of time	$\text{Poi}(\lambda)$	(next week) Interval of time to first success

Geometric RV

Consider an experiment: independent trials of $\text{Ber}(p)$ random variables.

def A **Geometric** random variable X is the # of trials until the first success.

$$X \sim \text{Geo}(p)$$

Support: $\{1, 2, \dots\}$

PMF

$$P(X = k) = (1 - p)^{k-1}p$$

Expectation

$$E[X] = \frac{1}{p}$$

Variance

$$\text{Var}(X) = \frac{1-p}{p^2}$$

Examples:

- Flipping a coin ($P(\text{heads}) = p$) until first heads appears
- Generate bits with $P(\text{bit} = 1) = p$ until first 1 generated

Negative Binomial RV

Consider an experiment: independent trials of $\text{Ber}(p)$ random variables.

def A **Negative Binomial** random variable X is the # of trials until r successes.


$X \sim \text{NegBin}(r, p)$	PMF	$P(X = k) = \binom{k-1}{r-1} (1-p)^{k-r} p^r$
	Expectation	$E[X] = \frac{r}{p}$
Support: $\{r, r+1, \dots\}$	Variance	$\text{Var}(X) = \frac{r(1-p)}{p^2}$

Examples:

- Flipping a coin until r^{th} heads appears
- # of strings to hash into table until bucket 1 has r entries

$$\text{Geo}(p) = \text{NegBin}(1, p)$$

Grid of random variables




Catching Pokemon

Wild Pokemon are captured by throwing Pokeballs at them.

- Each ball has probability $p = 0.1$ of capturing the Pokemon.
- Each ball is an independent trial.

What is the probability that you catch the Pokemon on the 5th try?



1. Define events/ RVs & state goal

$X \sim$ some distribution

Want: $P(X = 5)$

2. Solve

- A. $X \sim \text{Bin}(5, 0.1)$
- B. $X \sim \text{Poi}(0.5)$
- C. $X \sim \text{NegBin}(5, 0.1)$
- D. $X \sim \text{NegBin}(1, 0.1)$
- E. $X \sim \text{Geo}(0.1)$




Catching Pokemon

$$X \sim \text{Geo}(p) \quad p(k) = (1 - p)^{k-1}p$$

Wild Pokemon are captured by throwing Pokeballs at them.

- Each ball has probability $p = 0.1$ of capturing the Pokemon.
- Each ball is an independent trial.

What is the probability that you catch the Pokemon on the 5th try?



1. Define events/ RVs & state goal

$X \sim$ some distribution

Want: $P(X = 5)$


2. Solve

- A. $X \sim \text{Bin}(5, 0.1)$
- B. $X \sim \text{Poi}(0.5)$
- C. $X \sim \text{NegBin}(5, 0.1)$
- D. $X \sim \text{NegBin}(1, 0.1)$
- E. $X \sim \text{Geo}(0.1)$

CS109 Learning Goal: Use new RVs

Let's say you are learning about servers and networks.

You read about the M/D/1 queue:



"The service time busy period is distributed as a Borel with parameter $\mu = 0.2$."

Goal: You can recognize terminology and understand experiment setup.

Wikipedia page for Borel distribution. The page title is "Borel distribution". It includes a table of parameters, support, pmf, mean, and variance. The pmf is given as $e^{-\mu n} (\mu n)^{n-1} / n!$. The mean is $1/(1-\mu)$ and the variance is $\mu / (1-\mu)^3$. The page also includes a definition and a list of contents.

Borel distribution	
Parameters	$\mu \in [0, 1]$
Support	$n \in \{1, 2, 3, \dots\}$
pmf	$\frac{e^{-\mu n} (\mu n)^{n-1}}{n!}$
Mean	$\frac{1}{1-\mu}$
Variance	$\frac{\mu}{(1-\mu)^3}$

Definition [edit]

A discrete random variable X is said to have a Borel distribution^{[1][2]} with parameter $\mu \in [0, 1]$ if the probability mass function of X is given by

$$P_{\mu}(n) = \Pr(X = n) = \frac{e^{-\mu n} (\mu n)^{n-1}}{n!}$$

for $n = 1, 2, 3, \dots$



Exercises



The hardest part of is almost always deciding what you're modeling and what random variable to use.

Kickboxing with RVs

How might you model the following?

1. # of snapchats you receive in a day
2. # of children born to the same parents until the first one with green eyes
3. If stock went up (1) or down (0) in a day
4. # of probability problems you try until you get 5 correct (if you are randomly correct)
5. # of years since between now and 2050 with more than 6 Atlantic hurricanes

Choose from:

- A. $\text{Ber}(p)$
- B. $\text{Bin}(n, p)$
- C. $\text{Poi}(\lambda)$
- D. $\text{Geo}(p)$
- E. $\text{NegBin}(r, p)$



Kickboxing with RVs

How might you model the following?

1. # of snapchats you receive in a day
2. # of children born to the same parents until the first one with green eyes
3. If stock went up (1) or down (0) in a day
4. # of probability problems you try until you get 5 correct (if you are randomly correct)
5. # of years since between now and 2050 with more than 6 Atlantic hurricanes

Note: These exercises are designed to build intuition; in a problem statement, you'll generally be given more detail.

Choose from:

A. Ber(p)	C. Poi(λ)
B. Bin(n, p)	D. Geo(p)
	E. NegBin(r, p)

- C. Poi(λ)
- D. Geo(p) or E. NegBin(1, p)
- A. Ber(p) or B. Bin(1, p)
- E. NegBin($r = 5, p$)
- B. Bin($n = 27, p$), where $p = P(\geq 6 \text{ hurricanes in a year})$ calculated from C. Poi(λ)

Poisson Random Variable

Review

$$X \sim \text{Poi}(\lambda)$$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation

$$E[X] = \lambda$$

Support: $\{0, 1, 2, \dots\}$

Variance

$$\text{Var}(X) = \lambda$$

In CS109, a Poisson RV $X \sim \text{Poi}(\lambda)$ most often models

1. # of successes in a fixed interval of time, where successes are independent
 $\lambda = E[X]$, average success/interval

1. Web server load

$$\begin{array}{l} X \sim \text{Poi}(\lambda) \\ E[X] = \lambda \end{array} \quad p(k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Consider requests to a web server in 1 second.

- In the past, server load averages 2 hits/second, where requests arrive independently.
- Let $X = \#$ requests the server receives in a second.

What is $P(X < 5)$?

Define RVs

Solve

Poisson Random Variable


$X \sim \text{Poi}(\lambda)$	PMF	$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$
	Expectation	$E[X] = \lambda$
	Variance	$\text{Var}(X) = \lambda$
Support: $\{0, 1, 2, \dots\}$		

In CS109, a Poisson RV $X \sim \text{Poi}(\lambda)$ most often models

1. # of successes in a fixed time interval, where successes are independent
 $\lambda = E[X]$, average success/interval
2. Approximation of $Y \sim \text{Bin}(n, p)$ where n is large and p is small.
 $\lambda = E[Y] = np$

Approximation works well even when trials not entirely independent.

2. DNA



All the movies, images, emails and other digital data from more than 600 smartphones (10,000 GB) can be stored in the faint pink smear of DNA at the end of this test tube.

What is the probability that DNA storage stays uncorrupted?

2. DNA

What is the probability that DNA storage stays uncorrupted?


- In DNA (and real networks), we store large strings.
- Let string length be long, e.g., $n \approx 10^4$
- Probability of corruption of each base pair is very small, e.g., $p = 10^{-6}$
- Let $X = \#$ of corruptions.

What is $P(\text{DNA storage is uncorrupted}) = P(X = 0)$?

1. Approach 1:

$$X \sim \text{Bin}(n = 10^4, p = 10^{-6})$$

$$P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$

unwieldy!  $= \binom{10^4}{0} 10^{-6 \cdot 0} (1 - 10^{-6})^{10^4 - 0}$
 ≈ 0.990049829


2. Approach 2:

$$X \sim \text{Poi}(\lambda = 10^4 \cdot 10^{-6} = 0.01)$$

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!} = e^{-0.01} \frac{0.01^0}{0!}$$

$$= e^{-0.01}$$

$$\approx 0.990049834$$

a good approximation! 

When is a Poisson approximation appropriate?

$$P(X = k) = \lim_{n \rightarrow \infty} \binom{n}{k} \left(\frac{\lambda}{n}\right)^k \left(1 - \frac{\lambda}{n}\right)^{n-k} = \dots$$

Def natural
exponent

$$= \lim_{n \rightarrow \infty} \frac{n!}{n^k (n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

Expand

$$= \lim_{n \rightarrow \infty} \frac{n(n-1) \cdots (n-k+1)}{n^k} \frac{(n-k)!}{(n-k)!} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{\left(1 - \frac{\lambda}{n}\right)^k}$$

Limit analysis

$$= \lim_{n \rightarrow \infty} \frac{n^k}{n^k} \frac{\lambda^k}{k!} \frac{e^{-\lambda}}{1}$$

Simplify

$$= \frac{\lambda^k}{k!} e^{-\lambda}$$

Under which conditions will
 $X \sim \text{Bin}(n, p)$ behave like
 $\text{Poi}(\lambda)$, where $\lambda = np$?

- A. Large n , large p
- B. Small n , small p
- C. Large n , small p
- D. Small n , large p
- E. Other



Poisson approximation

$$X \sim \text{Poi}(\lambda)$$
$$E[X] = \lambda$$

$$Y \sim \text{Bin}(n, p)$$
$$E[Y] = np$$


Poisson approximates Binomial when n is large, p is small, and $\lambda = np$ is "moderate".

Different interpretations of "moderate":

- $n > 20$ and $p < 0.05$
- $n > 100$ and $p < 0.1$

Poisson is Binomial in the limit:

- $\lambda = np$, where $n \rightarrow \infty, p \rightarrow 0$



Poisson Random Variable

Consider an experiment that lasts a fixed interval of time.

def A **Poisson** random variable X is the number of occurrences over the experiment duration.

$$X \sim \text{Poi}(\lambda)$$

Support: $\{0, 1, 2, \dots\}$

PMF

$$P(X = k) = e^{-\lambda} \frac{\lambda^k}{k!}$$

Expectation $E[X] = \lambda$

Variance $\text{Var}(X) = \lambda$

Examples:

- # earthquakes per year
- # server hits per second
- # of emails per day

Time to show intuition for why expectation == variance!

Properties of $\text{Poi}(\lambda)$ with the Poisson paradigm

Recall the Binomial:

$Y \sim \text{Bin}(n, p)$	Expectation	$E[Y] = np$
	Variance	$\text{Var}(Y) = np(1 - p)$

Consider $X \sim \text{Poi}(\lambda)$, where $\lambda = np$ ($n \rightarrow \infty, p \rightarrow 0$):

$X \sim \text{Poi}(\lambda)$	Expectation	$E[X] = \lambda$
	Variance	$\text{Var}(X) = \lambda$

Proof:


$$E[X] = np = \lambda$$
$$\text{Var}(X) = np(1 - p) \rightarrow \lambda(1 - 0) = \lambda$$



Poisson Approximation, approximately

Poisson can still provide a **good approximation of the Binomial**, even when assumptions are "mildly" violated.

You can apply the Poisson approximation when:

- "Successes" in trials are not entirely independent e.g., # entries in each bucket in large hash table. 
- Probability of "success" in each trial varies (slightly), like a **small relative change** in a very small p e.g., average # requests to web server/sec may fluctuate slightly due to load on network

We won't explore this too much, but we want you to know about it anyway.

Can these Binomial RVs be approximated?




Poisson approximates Binomial when n is large, p is small, and $\lambda = np$ is "moderate".

Different interpretations of "moderate":

- $n > 20$ and $p < 0.05$
- $n > 100$ and $p < 0.1$

Poisson is Binomial in the limit:

- $\lambda = np$, where $n \rightarrow \infty, p \rightarrow 0$



Can these Binomial RVs be approximated?

Poisson approximates Binomial when n is large, p is small, and $\lambda = np$ is "moderate".

Different interpretations of "moderate":

- $n > 20$ and $p < 0.05$
- $n > 100$ and $p < 0.1$

Poisson is Binomial in the limit:

- $\lambda = np$, where $n \rightarrow \infty, p \rightarrow 0$

