

(1) (10 points) Find bases of the null space and the column space of the matrix

$$A = \begin{pmatrix} 1 & 2 & 0 & 1 & 2 \\ 1 & 2 & 0 & 2 & 3 \\ 1 & 2 & 0 & 3 & 4 \\ 1 & 2 & 0 & 4 & 5 \end{pmatrix}$$

Find $\text{rref}(A)$.

$$\begin{array}{l} R2-R1 \\ R3-R1 \\ R4-R1 \end{array} \begin{pmatrix} 1 & 2 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 2 & 2 \\ 0 & 0 & 0 & 3 & 3 \end{pmatrix} \rightsquigarrow \begin{array}{l} R1-R2 \\ R3-R2 \\ R4-R2 \end{array} \begin{pmatrix} 1 & 2 & 0 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix} = \text{rref}(A)$$

x_1 x_2 x_3 x_4 x_5
 \uparrow \swarrow \uparrow \swarrow
 pivot free pivot free

The nullspace is vectors $\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = x_2 \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix} + x_3 \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix} + x_5 \begin{bmatrix} -1 \\ 0 \\ 0 \\ -1 \\ 1 \end{bmatrix}$,

so a basis for $N(A)$ is $\left\{ \begin{bmatrix} -2 \\ 1 \\ 0 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 0 \\ 0 \\ -1 \\ 1 \end{bmatrix} \right\}$.

A basis for $C(A)$ corresponds to pivot vectors of $\text{rref}(A)$ back in A — these are the linearly independent columns of A :

$$\left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 2 \\ 3 \\ 4 \end{bmatrix} \right\}$$

(2) (8 points) What condition(s) must b_1, b_2, b_3 and b_4 satisfy so that the following system has a solution?

$$x - 3y = b_1$$

$$3x + y = b_2$$

$$x + 7y = b_3$$

$$2x + 4y = b_4$$

The augmented matrix is

$$\left[\begin{array}{cc|c} 1 & -3 & b_1 \\ 3 & 1 & b_2 \\ 1 & 7 & b_3 \\ 2 & 4 & b_4 \end{array} \right] \rightsquigarrow \left[\begin{array}{cc|c} 1 & -3 & b_1 \\ 0 & 10 & b_2 - 3b_1 \\ 0 & 10 & b_3 - b_1 \\ 0 & 10 & b_4 - 2b_1 \end{array} \right] \begin{array}{l} \\ R2 - 3R1 \\ R3 - R1 \\ R4 - 2R1 \end{array}$$

$$\rightsquigarrow \left[\begin{array}{cc|c} 1 & 0 & b_1 + 3(b_2 - 3b_1)/10 \\ 0 & 1 & (b_2 - 3b_1)/10 \\ 0 & 0 & b_3 - b_1 - b_2 + 3b_1 \\ 0 & 0 & b_4 - 2b_1 - b_2 + 3b_1 \end{array} \right] \begin{array}{l} R1 + 3R2 \\ \text{new} \\ \\ \\ R4 - R2 \end{array}$$

For the system to have a solution, it must be consistent, that is

$$0 = b_3 - b_1 - b_2 + 3b_1$$

$$0 = b_4 - 2b_1 - b_2 + 3b_1.$$

Simplifying, b_1, b_2, b_3, b_4 must satisfy

$$\begin{cases} 0 = 2b_1 - b_2 + b_3 \\ 0 = b_1 - b_2 + b_4. \end{cases}$$

- (5) (10 points) A box containing pennies, nickels and dimes contains 13 coins altogether, with a total value of 83 cents. How many coins of each type are in the box?

To make 83¢, there can be

3

8

13

pennies, but with 8 and 13, you couldn't make 83¢ with only 5 or 0 more coins. So there are 3 pennies.

Let $n = \#$ nickels
 $d = \#$ dimes.

$$\text{Then } n + d = 10$$

$$5n + 10d = 80$$

$$\begin{array}{c} \downarrow \\ \left[\begin{array}{cc|c} 1 & 1 & 10 \\ 5 & 10 & 80 \end{array} \right] \rightsquigarrow R_2 - 5R_1 \left[\begin{array}{cc|c} 1 & 1 & 10 \\ 0 & 5 & 30 \end{array} \right] \end{array}$$

$$\rightsquigarrow \begin{array}{l} R_1 - R_2 \\ 1/6 R_2 \end{array} \left[\begin{array}{cc|c} 1 & 0 & 4 \\ 0 & 1 & 6 \end{array} \right] \rightsquigarrow \begin{array}{l} 4 \text{ nickels} \\ 6 \text{ dimes} \\ 3 \text{ pennies} \end{array}$$

(6) (17 points) Let

$$V = \text{span} \left\{ \begin{bmatrix} -2 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -1 \\ 1 \\ 0 \\ 1 \end{bmatrix} \right\}, \quad u = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix}, \quad \vec{v}_1 = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \quad \vec{v}_2 = \begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix}.$$

(a) Show that \vec{v}_1 and \vec{v}_2 belong to the orthogonal complement V^\perp of V .

$$\begin{bmatrix} -2 \\ 1 \\ 1 \\ 0 \end{bmatrix} \cdot \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} = -2 \cdot 1 + 1 \cdot 1 + 1 \cdot 1 = 0$$

$$\text{and } \begin{bmatrix} -1 \\ 1 \\ 0 \\ 1 \end{bmatrix} \cdot \vec{v}_1 = -1 \cdot 1 + 1 \cdot 0 + 0 \cdot 2 + 1 \cdot 1 = 0 \Rightarrow \vec{v}_1 \text{ is in } V^\perp$$

$$\text{Similarly } \begin{bmatrix} -2 \\ 1 \\ 1 \\ 0 \end{bmatrix} \cdot \vec{v}_2 = -2 \cdot 1 + 1 \cdot 0 + 1 \cdot 2 + 0 \cdot 1 = 0 \quad \text{and} \quad \begin{bmatrix} -1 \\ 1 \\ 0 \\ 1 \end{bmatrix} \cdot \vec{v}_2 = -1 \cdot 1 + 1 \cdot 0 + 0 \cdot 2 + 1 \cdot 1 = 0$$

$$\Rightarrow \vec{v}_2 \text{ is in } V^\perp$$

(b) Is $\{\vec{v}_1, \vec{v}_2\}$ a basis of V^\perp ? Explain why or why not.

Yes.

Since V is a 2 dimensional subspace of \mathbb{R}^4 , the dimension of V^\perp is $4 - 2 = 2$.

\vec{v}_1 and \vec{v}_2 are 2 linearly independent vectors in V^\perp , so they also span V^\perp .

Thus they form a basis for V^\perp .

(c) Find an orthonormal basis of V^\perp .

Gram-Schmidt

$$\vec{w}_1 = \frac{\vec{v}_1}{\|\vec{v}_1\|} = \frac{1}{\sqrt{1^2+1^2+1^2+0^2}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}$$

$$\begin{aligned} \vec{y}_2 &= \vec{v}_2 - \text{proj}_{V_1} \vec{v}_2 = \vec{v}_2 - (\vec{w}_1 \cdot \vec{v}_2) \vec{w}_1 \\ &= \begin{bmatrix} 1 \\ 0 \\ 2 \\ 1 \end{bmatrix} - \underbrace{\frac{1}{\sqrt{3}}(1 \cdot 1 + 1 \cdot 0 + 1 \cdot 2 + 0 \cdot 1)}_{\frac{3}{3}=1} \left(\frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} \right) \\ &= \begin{bmatrix} 0 \\ -1 \\ 1 \\ 1 \end{bmatrix} \end{aligned}$$

$$\vec{w}_2 = \frac{\vec{y}_2}{\|\vec{y}_2\|} = \frac{1}{\sqrt{3}} \begin{bmatrix} 0 \\ -1 \\ 1 \\ 1 \end{bmatrix}$$

$$\left\{ \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix}, \frac{1}{\sqrt{3}} \begin{bmatrix} 0 \\ -1 \\ 1 \\ 1 \end{bmatrix} \right\}$$

(d) Find the orthogonal projection of u on V .

Use that $\text{proj}_V \vec{u} + \text{proj}_{V^\perp} \vec{u} = I_4(\vec{u}) = \vec{u}$

$$\text{proj}_{V^\perp} \vec{u} = (\vec{u} \cdot \vec{w}_1) \vec{w}_1 + (\vec{u} \cdot \vec{w}_2) \vec{w}_2$$

$$= \underbrace{\left(\frac{1}{\sqrt{3}}(1 \cdot 1 + 1 \cdot 1 + 1 \cdot 1 + 1 \cdot 0) \right)}_{=1} \frac{1}{\sqrt{3}} \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} + \underbrace{\left(\frac{1}{\sqrt{3}}(1 \cdot 0 + 1 \cdot -1 + 1 \cdot 1 + 1 \cdot 1) \right)}_{=1/3} \frac{1}{\sqrt{3}} \begin{bmatrix} 0 \\ -1 \\ 1 \\ 1 \end{bmatrix}$$

$$= \begin{bmatrix} 1 \\ 1 \\ 1 \\ 0 \end{bmatrix} + \frac{1}{3} \begin{bmatrix} 0 \\ -1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2/3 \\ 4/3 \\ 1/3 \end{bmatrix}$$

$$\text{So } \text{proj}_V \vec{u} = \vec{u} - \text{proj}_{V^\perp} \vec{u} = \begin{bmatrix} 1 \\ 1 \\ 1 \\ 1 \end{bmatrix} - \begin{bmatrix} 1 \\ 2/3 \\ 4/3 \\ 1/3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1/3 \\ -1/3 \\ 2/3 \end{bmatrix}$$

(7) (10 points) Let $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be projection onto the plane P that passes through $\vec{0}$ and is orthogonal to the line spanned by $\begin{bmatrix} 1 \\ 0 \\ 9 \end{bmatrix}$.

(a) Find an eigenbasis for T .

T has eigenvalues 0 and 1
 with eigenspace the line spanned by $\begin{bmatrix} 1 \\ 0 \\ 9 \end{bmatrix}$ and the plane P

$$1x + 0y + 9z = 0$$

$$\downarrow \\ x = -9z$$

So an eigenbasis is

$$\left\{ \begin{bmatrix} 1 \\ 0 \\ 9 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -9 \\ 0 \\ 1 \end{bmatrix} \right\}$$

$$P = \text{span} \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} = y \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + z \begin{bmatrix} -9 \\ 0 \\ 1 \end{bmatrix} \right\}$$

(b) Write down a matrix in standard coordinates which represents T . You can express your matrix as a product of matrices and inverses of matrices.

$$A = \begin{bmatrix} 0 & -9 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$A^T A = \begin{bmatrix} 0 & 1 & 0 \\ -9 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 & -9 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 82 \end{bmatrix}$$

matrix for T is

$$A(A^T A)^{-1} A^T = \begin{bmatrix} 0 & -9 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 82 \end{bmatrix}^{-1} \begin{bmatrix} 0 & 1 & 0 \\ -9 & 0 & 1 \end{bmatrix}$$

Alternative: P^\perp is $\text{span} \left(\begin{bmatrix} 1 \\ 0 \\ 9 \end{bmatrix} \right)$. Use $\text{proj}_P = I_3 - \text{proj}_{P^\perp}$

Let $B = \begin{bmatrix} 1 \\ 0 \\ 9 \end{bmatrix}$. Then a matrix for proj_{P^\perp} is

$$B(B^T B)^{-1} B^T = \begin{bmatrix} 1 \\ 0 \\ 9 \end{bmatrix} [82]^{-1} [10 \ 9] \text{ so } \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} - \begin{bmatrix} 1 \\ 0 \\ 9 \end{bmatrix} [82]^{-1} [10 \ 9]$$

- (8) (15 points) Globo-tech Marketing monitors the dollars spent each year by its customers on apples and oranges. With $a(k)$ representing the number of dollars spent (in millions) on apples in year k , and $o(k)$ the number of dollars spent (in millions) on oranges in year k , they determine that

$$a(k+1) = \frac{2}{10}a(k) + \frac{4}{10}o(k)$$

$$o(k+1) = \frac{8}{10}a(k) + \frac{6}{10}o(k)$$

We shall write $\vec{v}_k = \begin{bmatrix} a(k) \\ o(k) \end{bmatrix}$.

- (a) Find a matrix A so that $A\vec{v}_k = \vec{v}_{k+1}$. Notice that this will imply $A^k\vec{v}_0 = \vec{v}_k$.

$$\begin{bmatrix} 2/10 & 4/10 \\ 8/10 & 6/10 \end{bmatrix}$$

- (b) Find the eigenvalues of A , and for each eigenvalue find a basis for the corresponding eigenspace.

$$\lambda I - A = \begin{bmatrix} \lambda - 2/10 & -4/10 \\ -8/10 & \lambda - 6/10 \end{bmatrix}$$

$$\begin{aligned} \det(\lambda I - A) &= (\lambda - 2/10)(\lambda - 6/10) - \frac{32}{100} \\ &= \lambda^2 - \frac{8}{10}\lambda + \frac{12}{100} - \frac{32}{100} \\ &= \lambda^2 - \frac{4}{5}\lambda - \frac{1}{5} = 0 \\ &(\lambda - 1)(\lambda + 1/5) = 0 \end{aligned}$$

Each entry is ≥ 0 and sum of entries in column is 1.
Note A is a Markov matrix, so 1 is an eigenvalue.

Eigenvalues $\lambda = 1, -1/5$

$$E_1 = N(I - A)$$

$$\begin{bmatrix} 4/5 & -2/5 \\ -4/5 & 2/5 \end{bmatrix} \xrightarrow{R_2 + R_1} \begin{bmatrix} 4/5 & -2/5 \\ 0 & 0 \end{bmatrix}$$

$$\xrightarrow{5/4 \cdot R_1} \begin{bmatrix} 1 & -1/2 \\ 0 & 0 \end{bmatrix} \quad \text{implies} \quad \begin{aligned} x_1 - 1/2 x_2 &= 0 \\ \downarrow \\ x_1 &= 1/2 x_2 \end{aligned}$$

$$E_1 = \left\{ \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_2 \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} \right\} \\ = \text{span} \left\{ \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} \right\}$$

$$E_{-1/5} = N(-1/5 I - A)$$

$$\begin{bmatrix} -2/5 & -2/5 \\ -4/5 & -4/5 \end{bmatrix} \xrightarrow{R_2 - 2R_1} \begin{bmatrix} -2/5 & -2/5 \\ 0 & 0 \end{bmatrix}$$

$$\xrightarrow{-5/2 R_1} \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix} \quad \text{implies} \quad \begin{aligned} x_1 + x_2 &= 0 \\ \downarrow \\ x_1 &= -x_2 \end{aligned}$$

$$E_{-1/5} = \left\{ \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = x_2 \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\} \\ = \text{span} \left\{ \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right\}$$

(c) Express $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$ as a linear combination of the eigenvectors you just computed.

$$\begin{bmatrix} 2 \\ 1 \end{bmatrix} = c_1 \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} + c_2 \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

Find c_1 and c_2 .

$$2 = \frac{1}{2}c_1 - c_2$$

$$1 = c_1 + c_2 \rightarrow c_2 = 1 - c_1$$

$$\rightarrow 2 = \frac{1}{2}c_1 - 1 + c_1 \rightarrow 3 = \frac{3}{2}c_1$$

↓

$$c_1 = 2$$

↓

$$c_2 = -1$$

$$\text{So } \begin{bmatrix} 2 \\ 1 \end{bmatrix} = 2 \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} - \begin{bmatrix} -1 \\ 1 \end{bmatrix}.$$

(d) Suppose that $\vec{v}_0 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$. Using your answers from above, what is a good estimate for the number of dollars (in millions) spent on apples in year 100? What about dollars (in millions) spent on oranges in year 100?

$$A^{100} \vec{v}_0 = A^{100} \left[2 \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} - \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right]$$

$$= 2 A^{100} \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} - A^{100} \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

$$= 2 \begin{bmatrix} 1/2 \\ 1 \end{bmatrix} - \frac{1}{5^{100}} \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

≈ 0

$$\approx \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

1 million on apples
2 million on oranges

- (9) (8 points) Show that if A is an $n \times n$ matrix then there exist scalars c_0, \dots, c_n —not all zero—so that

$$\det(c_0 I_n + c_1 A + c_2 A^2 + \dots + c_n A^n) = 0.$$

(Hint: For a vector \vec{v} , what can you say about linear dependence of the collection $\vec{v}, A\vec{v}, \dots, A^n \vec{v}$? Why might this help you?)

If \vec{v} is a nonzero vector in \mathbb{R}^n ,
 $\vec{v}, A\vec{v}, \dots, A^n \vec{v}$ is a collection
of $(n+1)$ vectors in \mathbb{R}^n ,
so it must be linearly dependent.

Thus there are scalars c_0, \dots, c_n , not
all zero, so that

$$c_0 \vec{v} + c_1 A\vec{v} + \dots + c_n A^n \vec{v} = 0.$$

$$\parallel$$

$$(c_0 I_n + c_1 A + \dots + c_n A^n) \vec{v}$$

This implies \vec{v} is a nonzero vector
in the nullspace of $c_0 I_n + c_1 A + \dots + c_n A^n$,
i.e. $N(c_0 I_n + \dots + c_n A^n) \neq \{0\}$.

$$\Rightarrow \det(c_0 I_n + c_1 A + \dots + c_n A^n) = 0.$$

(10) (5 points) Does there exist a constant c such that

$$f(x,y) = \begin{cases} \frac{(x+y)^2}{x^2+y^2} & \text{if } (x,y) \neq (0,0) \\ c & \text{if } (x,y) = (0,0) \end{cases}$$

is continuous? Why or why not?

NO

$$\lim_{\substack{(x,y) \rightarrow (0,0) \\ \text{along} \\ y=mx}} \frac{(x+y)^2}{x^2+y^2} = \lim_{x \rightarrow 0} \frac{(x+mx)^2}{x^2+m^2x^2}$$

$$= \lim_{x \rightarrow 0} \frac{(1+m)^2 x^2}{(1+m^2) x^2} = \frac{(1+m)^2}{1+m^2}. \quad \text{Since the}$$

limit of $f(x,y)$ approaching $(0,0)$ along different lines $y=mx$ is different depending on m , the limit does not exist. So no value of c would make f continuous.

(11) (5 points) Let S be the surface in \mathbb{R}^3 defined by

$$x^2 + \frac{y^2}{4} - z^2 = 1.$$

What is the tangent plane to this surface at the point $(1,2,1)$?

$$f(x,y,z) = x^2 + \frac{y^2}{4} - z^2$$

$$\nabla f(x,y,z) = (2x, \frac{1}{2}y, -2z)$$

$$\nabla f(1,2,1) = (2, 1, -2) \quad \text{This is the normal vector to the plane.}$$

$$2(x-1) + 1(y-2) - 2(z-1) = 0$$

(12) (12 points) Consider the function $f(x, y) = x^2/y^4$.


(a) Carefully draw the level curve passing through the point $(1, -1)$. On this graph, draw the gradient of the function f at $(1, -1)$.

$$\text{at } (1, -1), f(x, y) = f(1, -1) = \frac{1^2}{(-1)^4} = 1.$$

$$\text{Thus the level curve is } f(x, y) = \frac{x^2}{y^4} = 1$$

$$\Rightarrow x^2 = y^4$$

$$\Rightarrow x = \pm y^2$$



$$\nabla f(x, y) = \begin{bmatrix} 2x/y^4 \\ -4x^2/y^5 \end{bmatrix}$$

$$\nabla f(1, -1) = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

(b) Compute the directional derivative of f at the point $(1, -1)$ in the direction $\vec{u} = \begin{bmatrix} 4/5 \\ 3/5 \end{bmatrix}$.

$$D_{\vec{u}} f = \nabla f(1, -1) \cdot \vec{u}$$

$$= \begin{bmatrix} 2 \\ 4 \end{bmatrix} \cdot \begin{bmatrix} 4/5 \\ 3/5 \end{bmatrix}$$

$$= 8/5 + 12/5 = 20/5 = \textcircled{4}$$

(c) Suppose that $f(x, y)$ gives the height of a mountain above (x, y) , and suppose further that you are stuck on the mountain at position $(1, -1, f(1, -1))$. In what direction $\begin{pmatrix} \Delta x \\ \Delta y \end{pmatrix}$ should you take your first step if you want to descend the mountain as quickly as possible?

$\nabla f(1, -1)$ is the direction of steepest ascent
 $-\nabla f(1, -1)$ " " descent.

$$\text{So walk in } \begin{pmatrix} \Delta x \\ \Delta y \end{pmatrix} = \begin{pmatrix} -2 \\ -4 \end{pmatrix}.$$

(13) (10 points) Consider the function

$$f(x, y, z) = \sqrt{\ln(e^{2x}yz^3)}$$

(a) Write down the first order Taylor polynomial centered at the point $(2, 1, 1)$.

$$p_1(x, y, z) = f(2, 1, 1) + [f_x \ f_y \ f_z] \begin{bmatrix} x-2 \\ y-1 \\ z-1 \end{bmatrix}$$

$$f(2, 1, 1) = \sqrt{\ln(e^4)} = \sqrt{4\ln e} = \sqrt{4} = 2$$

$$f_x = \frac{1}{2\sqrt{\ln(e^{2x}yz^3)}} \cdot \frac{1}{e^{2x}yz^3} \cdot \cancel{2e^{2x}}y z^3 \frac{at}{(2,1,1)} \frac{1}{\sqrt{\ln(e^4)}} = \frac{1}{2}$$

$$f_y = \frac{1}{2\sqrt{\ln(e^{2x}yz^3)}} \cdot \frac{1}{e^{2x}yz^3} \cdot e^{2x}z^3 \frac{at}{(2,1,1)} \frac{1}{2\sqrt{\ln(e^4)}} = \frac{1}{4}$$

$$f_z = \frac{1}{2\sqrt{\ln(e^{2x}yz^3)}} \cdot \frac{1}{e^{2x}yz^3} \cdot e^{2x}y 3z^2 \frac{at}{(2,1,1)} \frac{3}{2} = \frac{3}{4}$$

$$p_1(x, y, z) = 2 + \frac{1}{2}(x-2) + \frac{1}{4}(y-1) + \frac{3}{4}(z-1)$$

(b) Find the approximate value of the number $\sqrt{\ln(e^{4.01}(.98)(1.03)^3)}$.

$$f(2.005, .98, 1.03)$$

$$\approx p_1(2.005, .98, 1.03)$$

$$= 2 + \frac{1}{2}(.005) + \frac{1}{4}(-.02) + \frac{3}{4}(.03)$$

$$= 2.02$$

(14) (10 points) Find all critical points of the function $2x^3 + 6xy + 3y^2$ and describe their nature.

Find crit pts

$$\frac{\partial f}{\partial x} = 6x^2 + 6y = 0$$

$$\frac{\partial f}{\partial y} = 6x + 6y = 0$$

$$y = -x$$

$$x^2 - x = 0 \rightarrow x=0 \rightarrow y=0$$

$$(x)(x-1) \rightarrow x=1 \rightarrow y=-1$$

Two critical points: $(0,0)$ and $(1,-1)$

Nature

$$H_f = \begin{bmatrix} 12x & 6 \\ 6 & 6 \end{bmatrix}$$

$$H_f(0,0) = \begin{bmatrix} 0 & 6 \\ 6 & 6 \end{bmatrix}$$

$$d_1 = 0$$

$$d_2 = 0 \cdot 6 - 6^2 = -36 < 0$$

↓

$(0,0)$ is a saddle point.

$$H_f(1,-1) = \begin{bmatrix} 12 & 6 \\ 6 & 6 \end{bmatrix}$$


$$d_1 = 12 > 0$$

$$d_2 = 12 \cdot 6 - 6^2 > 0$$

↓

$(1,-1)$ is a local minimum.

- (15) (10 points) Use calculus to find the point on the circle $(x-1)^2 + (y-2)^2 = 1$ which is nearest to the origin.



Minimize

$f(x, y) =$ distance squared to origin

$$f(x, y) = x^2 + y^2$$

constraint: $(x-1)^2 + (y-2)^2 = 1 = g(x, y)$

$$\begin{array}{l} \textcircled{x} \\ \textcircled{y} \\ \textcircled{g} \end{array} \left\{ \begin{array}{l} f_x - \lambda g_x = 0 \\ f_y - \lambda g_y = 0 \\ g = 1 \end{array} \right. \rightarrow \begin{array}{l} 2x - \lambda(2(x-1)) = 0 \rightarrow \lambda = \frac{x}{x-1} \\ 2y - \lambda(2(y-2)) = 0 \rightarrow \lambda = \frac{y}{y-2} \\ (x-1)^2 + (y-2)^2 = 1 \end{array}$$

$$\frac{x}{x-1} = \frac{y}{y-2} \Rightarrow xy - 2x = xy - y \Rightarrow y = 2x$$

Plug into \textcircled{g} : $(x-1)^2 + (2x-2)^2 = 1$

$$\begin{array}{l} \text{"} \\ (2(x-1))^2 \\ \text{"} \\ 4(x-1)^2 \end{array} \Rightarrow 5(x-1)^2 = 1$$

$$(x-1)^2 = \frac{1}{5}$$

$$x-1 = \pm \sqrt{\frac{1}{5}}$$

$$x = 1 \pm \sqrt{\frac{1}{5}}$$

critical points are

$$(1 - \sqrt{\frac{1}{5}}, 2 - 2\sqrt{\frac{1}{5}}) \rightarrow \text{distance}^2 = 5 - 10\sqrt{\frac{1}{5}} \text{ smaller}$$

$$(1 + \sqrt{\frac{1}{5}}, 2 + 2\sqrt{\frac{1}{5}}) \rightarrow \text{distance}^2 = 5 + 10\sqrt{\frac{1}{5}} \text{ larger}$$

So $(1 - \sqrt{\frac{1}{5}}, 2 - 2\sqrt{\frac{1}{5}})$ is nearest to the origin.

(Geometrically, a local min is a global min, and there is only one closest point.)