

## MIDTERM 2

- Complete the following problems. You may use any result from class you like, but if you cite a theorem be sure to verify the hypotheses are satisfied.
- This is a closed-book, closed-notes exam. No calculators or other electronic aids will be permitted.
- In order to receive full credit, please show all of your work and justify your answers. You do not need to simplify your answers unless specifically instructed to do so.
- If you need extra room, use the back sides of each page. If you must use extra paper, make sure to write your name on it and attach it to this exam. Do not unstaple or detach pages from this exam.
- Please sign the following:

“On my honor, I have neither given nor received any aid on this examination. I have furthermore abided by all other aspects of the honor code with respect to this examination.”

Name: \_\_\_\_\_

Signature: \_\_\_\_\_

<b>1</b>	10 pts	
<b>2</b>	10 pts	
<b>3</b>	13 pts	
<b>4</b>	15 pts	
<b>5</b>	10 pts	
<b>6</b>	12 pts	
<b>7</b>	10 pts	
<b>Total</b>	80 pts	

\_\_\_\_\_  
Circle your TA's name  
\_\_\_\_\_

Lan  
Oren  
Josh  
Peter  
Chad  
Leo  
Rob  
Nikola  
Jian

- (1) (10 point) Find the determinant of each of the following matrices. Show your work or justify your answer.

$$(a) \begin{pmatrix} 2 & 3 & 1 & 0 \\ 1 & 0 & 2 & 2 \\ 3 & 5 & 6 & 9 \\ 2 & 0 & 3 & 3 \end{pmatrix}$$

$$(b) \begin{pmatrix} 1 & 2 & 3 & 5 & 10 & 0 & 3 \\ 2 & 3 & 1 & 1 & 0 & 0 & 2 \\ 3 & 5 & 4 & 6 & 10 & 0 & 5 \\ 0 & 7 & 2 & 3 & 4 & 2 & 1 \\ 4 & 1 & 8 & 9 & 10 & 6 & 7 \\ 3 & 1 & 5 & 1 & 1 & 10 & 3 \\ 2 & 0 & 0 & 3 & 4 & 1 & 8 \end{pmatrix} \text{ (Hint: What is row 3 in terms of rows 1 and 2?)}$$

(2) (10 points) Suppose that  $V \subseteq \mathbb{R}^4$  is a subspace with basis

$$\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ 2 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 6 \\ 3 \\ 4 \\ 0 \end{bmatrix} \right\}.$$

(a) Use the Gram-Schmidt process and  $\mathcal{B}$  to produce an orthonormal basis for  $V$ .

(b) Compute the orthogonal projection of  $\vec{x} = \begin{bmatrix} 9 \\ 0 \\ 25 \\ 9 \end{bmatrix}$  onto  $V$ .

(3) (13 points) One can show that  $\mathcal{B} = \{\vec{v}_1, \vec{v}_2, \vec{v}_3, \vec{v}_4\}$  is a basis for  $\mathbb{R}^4$ , where

$$\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \\ 3 \\ 4 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} 5 \\ 6 \\ 7 \\ 8 \end{bmatrix}, \vec{v}_3 = \begin{bmatrix} 1 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \vec{v}_4 = \begin{bmatrix} 1 \\ 0 \\ 3 \\ 0 \end{bmatrix}.$$

(You don't have to prove this!)

(a) Give the matrix  $C$  which changes basis from  $\mathcal{B}$  to the standard basis. That is, find  $C$  so that

$$C[\vec{v}]_{\mathcal{B}} = [\vec{v}]_{\mathcal{S}},$$

where  $[\vec{v}]_{\mathcal{B}}$  is  $\vec{v}$  in  $\mathcal{B}$ -coordinates and  $[\vec{v}]_{\mathcal{S}}$  is  $\vec{v}$  in standard coordinates.

(b) Suppose that  $T: \mathbb{R}^4 \rightarrow \mathbb{R}^4$  is the linear transformation which satisfies

- $T(\vec{v}_1) = \vec{v}_1 + 2\vec{v}_2$ ,
- $T(\vec{v}_2) = \vec{v}_1 + \vec{v}_3$ ,
- $T(\vec{v}_3) = \vec{v}_1 + \vec{v}_2 - \vec{v}_4$ , and
- $T(\vec{v}_4) = \vec{0}$ .

Give the matrix for  $T$  in coordinates relative to the basis  $\mathcal{B}$ .

- (c) Give the matrix for  $T$  in coordinates relative to the standard basis. (You may express your answer as a product of matrices and their inverses without expanding out the products or computing the inverses).

- (4) (15 points) Let

$$A = \begin{pmatrix} 1 & 3 & 1 \\ 0 & 5 & 1 \\ 0 & -9 & -1 \end{pmatrix}.$$

- (a) Find all eigenvalues of  $A$ .

(b) For each eigenvalue, give a basis for the corresponding eigenspace.

(c) Is  $A$  diagonalizable? Be sure to explain your answer fully.

- (5) (10 points) Suppose that  $\mathcal{B} = \{\vec{v}_1, \dots, \vec{v}_n\}$  is an orthonormal basis for  $\mathbb{R}^n$ , and let  $Q$  be the (square) matrix

$$Q = \begin{pmatrix} | & & | \\ \vec{v}_1 & \cdots & \vec{v}_n \\ | & & | \end{pmatrix}.$$

- (a) What is  $Q^T Q$ ? Justify your answer.

- (b) Prove that  $\|Q\vec{x}\| = \|\vec{x}\|$ .

(6) (12 points) Let  $V$  be a  $k$ -dimensional subspace of  $\mathbb{R}^n$  for some  $0 < k < n$ , and let  $T$  be the linear transformation which projects vectors onto  $V$ .

(a) Prove that there exists a non-zero vector in  $V^\perp$ .

(b) Prove that 0 is an eigenvalue of  $T$ .

(c) Prove that 1 is an eigenvalue of  $T$ .

(7) (10 points)

(a) Let

$$f(x, y) = \begin{cases} |y|, & \text{if } |x| \leq |y| \\ |x|, & \text{if } |y| < |x|. \end{cases}$$

Carefully draw (and label) the level curves  $f(x, y) = 0$ ,  $f(x, y) = 1$  and  $f(x, y) = 4$ .

(b) Determine the value of the constant  $c$  so that

$$g(x, y) = \begin{cases} \frac{x^3 + xy^2 + 2x^2 + 2y^2}{x^2 + y^2} & \text{if } (x, y) \neq (0, 0) \\ c & \text{if } (x, y) = (0, 0) \end{cases}$$

is continuous. Be sure to justify your answer.