

Affine Point Processes and Portfolio Credit Risk

Eymen Errais

Creditex

Kay Giesecke*

Stanford University

Lisa R. Goldberg

MSCI Barra

May 23, 2006; this draft March 30, 2009[†]

Abstract

A stochastic model of correlated event timing is at the center of applications in many areas, including finance, insurance, reliability, queuing and health care. The event dependence structure often involves feedback loops along with the influence of exogenous stochastic risk factors. This paper analyzes a family of multivariate point process models of correlated event timing whose arrival intensity is driven by an affine jump diffusion. The components of an affine point process are self- and cross-exciting, and facilitate the description of complex event dependence structures. Ordinary differential equations characterize the transform of an affine point process and the probability distribution of an integer-valued affine point process. The moments of an affine point process take a closed form. This guarantees a high degree of computational tractability in applications. We illustrate this in the context of portfolio credit risk, where the correlation of corporate defaults is the main issue. We consider the valuation of securities exposed to correlated default risk, and demonstrate the significance of our results in a market calibration exercise that addresses the turmoil in the credit markets during the fall of 2008.

Key words: Self-exciting point process, affine jump diffusion, Hawkes process, transform, portfolio credit derivative, correlated default, index and tranche swap

*Corresponding author. Department of Management Science & Engineering, Stanford University, Stanford, CA 94305-4026, USA, Phone (650) 723 9265, Fax (650) 723 1614, email: giesecke@stanford.edu, web: www.stanford.edu/~giesecke.

[†]This paper was previously circulated under the title “Pricing Credit From the Top-Down With Affine Point Processes.” We thank Andrew Abrahams, Benjamin Armbruster, Damiano Brigo, Peter Carr, Valdo Durreleman, Andreas Eckner, Bjorn Flesaker, Igor Halperin, Steven Hutt, Peter Jäckel, Rajnish Kamat, Andrei Lopatin, Woujiang Lou, Timur Misirpashaev, George Papanicolaou, Vijay Poduri, David Soronow, Tom Stone, Pascal Tomecek, Nicolas Victoir, Stefan Weber and Ulrich Zier for helpful discussions and comments. We are extremely grateful for excellent research assistance from Xiaowei Ding, Baeho Kim, Jack Kim and Kao-chih Syao.

1 Introduction

The collapse of Lehman Brothers on September 15, 2008 brought the financial system to the brink of a breakdown. The dramatic repercussions point to the existence of feedback phenomena that are channeled through the complex web of informational and contractual relationships in the economy. Lehman was an important node in a network of derivative traders. It had bought and sold default insurance on a large number of firms and was itself a reference entity in countless other insurance contracts. Its downfall triggered payments that forced some insurance sellers into default, leaving the corresponding protection buyers with losses. It also exposed the counterparties to the contracts Lehman itself had written.

This and related episodes motivate the design of models of correlated default timing that incorporate the feedback phenomena that plague credit markets. This paper develops a family of computationally tractable stochastic point processes that can capture event feedback, and therefore are well-suited to the analysis of credit risk. We study a multivariate *affine point process* whose event arrival intensity is driven by an affine jump diffusion process that represents a vector of stochastic risk factors. The variation of the factors generates diffusive and jump volatility of conditional event arrival rates. The point process itself can be a risk factor so that the future evolution of the point process is influenced by the timing of past events and their marks, for example the financial loss they caused. This *self-exciting* feature takes account of the direct impact of events, and generates a dependence structure between arrival rates and event marks. It also allows us to specify a multivariate point process whose component processes interact through cross-excitation. The resulting models significantly generalize the Poisson, birth and Hawkes processes, and have applications in many areas, some of which are discussed in Section 1.2.

Our stepping stone for the specification and analysis of affine point processes is the Hawkes process. The Hawkes process is, perhaps, the simplest self-exciting point process. Its intensity jumps in response to events, and tends toward a target level in the absence of an event. We develop the distributional properties of the Hawkes process based on the fact that the two-dimensional process consisting of a Hawkes process and its intensity is Markov. The structure of the associated infinitesimal generator leads to a Dynkin formula and closed expressions for the moments of the Hawkes intensity. We show that a transform of the Hawkes process, which includes the Laplace and Fourier transforms as special cases, satisfies a certain partial integral differential equation. The solution to that equation turns out to be an exponentially affine function of the intensity, whose coefficients satisfy a system of ordinary differential equations. Analysis of the transform leads to ordinary differential equations that characterize the probability distribution of a Hawkes process. We obtain closed formulas for the moments of the process.

The insight into the probabilistic structure of the Hawkes process leads us to propose an extension to a richer intensity process that is driven by an affine jump diffusion, and which may include the point process itself as a component. Our basic transform, distribution and moment characterization arguments extend to this family of affine point

processes. We obtain the joint transform of an affine point process and its driving risk factor process as an exponentially affine function of the current state, with coefficient functions satisfying ordinary differential equations. We show that, by constructing an extended state vector, this characterization can also be cast in terms of the transform results for affine jump diffusions developed by Duffie, Pan & Singleton (2000).

The transform, distribution and moment results generate computational tractability for a range of applications. To illustrate, we use an affine point process to model the cumulative loss due to default in a portfolio of firms. The jump times represent default times and the jump magnitudes represent the random losses at default. This formulation captures the impact of a default on the surviving names. It also incorporates the negative correlation between default and recovery rates. The transform results facilitate the valuation, hedging and calibration of a portfolio credit derivative, which is a security whose payoff is a specified function of the portfolio loss, and which provides insurance against default losses in the portfolio. An index swap, for example, pays any portfolio losses before the contract maturity. A tranche swap pays a slice of the portfolio loss specified by a lower and an upper attachment point. The values of these swaps are given in terms of an inverse transform. Our market calibration experiments, which are based on index and tranche price data observed before and after Lehman's collapse, indicate the empirical significance of the self-exciting property of the loss process.

1.1 Related credit risk literature

An affine point process facilitates a top-down approach to portfolio credit risk, in which the portfolio loss process is specified without reference to the constituent names. There are other examples of this approach in the literature. Arnsdorf & Halperin (2008) describe arrivals by a time-changed non-linear death process. Brigo, Pallavicini & Torresetti (2006) model events by a mixed Poisson process and its generalizations. Cont & Minca (2008) describe arrivals by a non-homogeneous Markov chain. Davis & Lo (2001) model events by a piece-wise deterministic Markov process. Ding, Giesecke & Tomecek (2006) consider a time-changed linear birth process as a model of arrivals. In Longstaff & Rajan (2008), defaults are driven by independent Poisson processes that model idiosyncratic, sector specific and economy-wide events. The family of loss processes proposed in this paper is distinct from, or in some special cases encompasses, the models introduced in these contributions. Our models feature multivariate, interacting point processes with affine structure, stochastic recoveries, correlated arrival and recovery rates, and semi-analytical transforms. Calibration experiments indicate the fit of an example specification to the market data during the recent market turmoil.

In an alternative bottom-up model design, the default intensity of each portfolio constituent are the primitives. The dependence among firms must be built into the single-name models. An example is a doubly-stochastic setting in which firm intensities are driven by common factors, as in Duffie & Gârleanu (2001), Eckner (2007), Kou & Peng

(2009), Papageorgiou & Sircar (2007) and others. Conditional on the factors, inter-arrival times are independent. A second example is a setting with interaction, in which a default triggers a response in the intensities of the surviving firms. In Giesecke & Weber (2006) and Jarrow & Yu (2001), this response is exogenous. In the “frailty” models of Collin-Dufresne, Goldstein & Helwege (2003), Duffie, Eckner, Horel & Saita (2008) and others, the response is the result of learning from default. A third example is a setting in which dependence between issuers is imposed by a copula function, as in Bassamboo, Juneja & Zeevi (2008), Chen & Glasserman (2008), Glasserman & Li (2005), Glasserman, Kang & Shahabuddin (2007) and others.

The top-down specification of the portfolio loss process generates computational advantages. However, an issue is the attribution of risk to the portfolio constituents. This comes up in the hedging of the constituent exposures associated with a position in a portfolio derivative. Constituent name hedging requires the sensitivities of the portfolio derivative price with respect to changes in the prices of the single-name derivatives referenced on the constituents. The sensitivities determine the amount of single-name protection on each portfolio constituent to be bought or sold in order to neutralize portfolio derivative price fluctuations due to small changes in the constituent risks. Giesecke & Goldberg (2005) develop a random thinning approach to estimate these hedges for a given portfolio loss process. They demonstrate the feasibility and effectiveness of this approach for the affine point processes analyzed in this paper.

While this article highlights derivatives valuation applications, affine point processes are also potential tools for the risk management of corporate debt portfolios, for which event feedback and the dependence between default and recovery rates are significant issues. For example, Das, Duffie, Kapadia & Saita (2007) test a bottom-up doubly-stochastic model whose features are similar to those of the models endorsed by the regulatory authorities to estimate portfolio credit risk. They find evidence of historical default clustering in excess of that implied by the model they tested. This suggests that doubly-stochastic models underestimate risk capital. A self-exciting affine point process model implies a more realistic degree of default clustering. It also accounts for the dependence between default and recovery rates, whose importance is emphasized in Basel II’s pillar I guidelines, see Basel Commission on Bank Regulation (2004).

1.2 Non-financial applications

Our results have applications in many other areas that involve the modeling of correlated event times. Affine point processes can be used to model the timing of *insurance* claims, and to price re-insurance instruments such as stop-loss contracts. Claim arrivals are governed by a stochastic intensity that is driven by exogenous risk factors. The self-exciting property can be important in situations where policy holders’ decision to file a claim is influenced by the actions of other policy holders. Affine point processes can also be used to model the *reliability* of component systems, and to estimate the probability of

system failure. Here, the self-exciting property describes a situation in which the failure of a component increases the stress on the other components, and may lead to a failure cascade. In *health care* applications, an affine point process can be used to describe the sequence of disease incidences, and to study the spread of the disease throughout a population. The self-exciting property is a key feature for the description of infectious diseases, and the cross-excitation property is useful for modeling the interaction between different infectious diseases. Other application areas include *queuing* and *traffic* systems.

1.3 Structure of this article

Section 2 analyzes the distributional properties of the Hawkes process, which serves as our stepping stone. We develop formulas for the transform of the Hawkes process, its moments and the moments of the Hawkes intensity process. Section 3 applies these results to the valuation of credit derivative securities. Market calibration experiments demonstrate the benefits of the self-exciting property, and the computational advantages of the transform approach. Section 4 extends the transform analysis to general affine point processes. It also provides further parametric examples of multivariate self- and cross-exciting point processes. Section 5 concludes. The appendix contains the proofs.

2 A basic self-exciting model

Consider a sequence of default stopping times $0 < T_1 < T_2 < \dots$ that are defined on a complete probability space (Ω, \mathcal{F}, P) with right-continuous and complete information filtration $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$. The nature of the probability measure P depends on the application. In risk management applications, P is the actual or statistical measure. In valuation applications, P is a risk-neutral pricing measure, relative to which the discounted price of a traded security is a martingale. The financial loss at T_n is given by a random variable $\ell_n \in \mathcal{F}_{T_n}$. The sequence (T_n, ℓ_n) generates a non-explosive default counting process N given by $N_t = \sum_{n \geq 1} 1_{\{T_n \leq t\}}$ and a loss point process L defined by $L_t = \sum_{n \geq 1} \ell_n 1_{\{T_n \leq t\}}$.

We propose to specify the processes N and L directly through a conditional arrival rate or intensity λ , and a distribution ν on $(0, \infty)$ for the loss ℓ_n at an event. We assume that the jump transform $\int e^{cz} d\nu(z)$ exists and is finite for complex c , and admits a finite derivative $\int ze^{cz} d\nu(z)$. The intensity follows a strictly positive stochastic process that describes the conditional mean default rate in the sense that $E(N_{t+\Delta} - N_t | \mathcal{F}_t) \approx \lambda_t \Delta$ for small $\Delta > 0$. This means that $N - \int_0^\cdot \lambda_s ds$ is a local martingale relative to P and \mathbb{F} . The process followed by λ completely determines the conditional distribution of N . We propose a family of models for λ whose features are empirically motivated, and which lead to a computationally tractable expression for a transform of N and L .




Figure 1: Annual defaults of Moody’s rated U.S. firms between January 1970 and November 17, 2008. Source: Moody’s Default Risk Service. The peak in 1970 represents a cluster of 24 railway defaults triggered by the collapse of Penn Central Railway on June 21, 1970. The fallout of the 1987 crash is indicated by the peak in the early nineties. The burst of the internet bubble caused many defaults during 2001-2002. From a trough in 2007, default rates increased significantly in 2008.

2.1 Empirical motivation

Empirical observation dictates the properties of the stochastic process followed by λ . Most importantly, λ must replicate the *clustering of defaults* seen in Figure 1, which is due to the dependence of the default times T_n . The dependence is a result of the sensitivity of firms to common economic risk factors that vary stochastically through time. It is also caused by the informational and contractual linkages between firms, which provide a channel for the propagation of financial distress from one firm to another. The existence of these feedback phenomena is indicated by the ripple effects associated with the default of Lehman Brothers on September 15, 2008, and is further empirically documented in Azizpour & Giesecke (2008), Collin-Dufresne et al. (2003), Duffie et al. (2008), Jorion & Zhang (2006), Longstaff (2008) and others. We propose to model the impact of a default on the other firms by ramping up the intensity at an event. This amounts to including the default process itself as a risk factor influencing the intensity.

Another important empirical feature is the negative correlation between default and recovery rates. During periods with elevated default rates, creditors tend to recover less at a default than during periods with relatively few defaults. While documented by Altman, Brady, Resti & Sironi (2005) and many others, this property has been largely ignored in a theoretical literature that is focused on modeling default timing. We propose to capture this property by modeling the magnitude of the response of the intensity to a default as a linear function of the realized loss at an event. The larger the loss, the larger the impact

of an event on the other firms, and the bigger the increase of the intensity at an event.

2.2 Specification

We propose a basic intensity model that incorporates the features described above. We suppose events arrive with intensity λ given by

$$\lambda_t = u(t) + \int_0^t h(t-s) dL_s. \quad (1)$$

The first to default intensity $u(t) = c + e^{-\kappa t}(\lambda_0 - c)$ is a deterministic function of time, $c > 0$, $\lambda_0 > 0$ and the impact of a loss on the intensity is governed by the function

$$h(v) = \delta e^{-\kappa v}, \quad v \geq 0, \quad (2)$$

with $\kappa \geq 0$ and $\delta \geq 0$. As illustrated in the sample path of (λ, L) shown in Figure 2, the randomness in the default rate is driven by two sources of uncertainty: the timing of events and the recovery at these events. At a default, the intensity ramps up by the realized loss scaled with the sensitivity parameter δ . The lower the recovery the higher the jump of the default rate. The impact of an event decays exponentially over time with rate κ . The reversion level is c . It follows that defaults are positively self-affecting, or *self-exciting*. Further, default and recovery rates are *negatively correlated*. Despite its parsimony, in Section 3.3 below we show that the basic model (1) is very effective at replicating the clustering of defaults implied by market prices.

The intensity λ governs the common event times of N and L . While the jumps of the default process N are unit-sized, the jumps of the loss process L are drawn from the distribution ν on $(0, \infty)$ of loss given default. The two dimensional process $J = (L, N)^\top$ is a Hawkes process, see Hawkes (1971) and Hawkes & Oakes (1974). Variants of the Hawkes process have been applied to a range of engineering problems, see Daley & Vere-Jones (2003). They have only recently been used in financial economics. For example, Bowsher (2007) fits different Hawkes models to security trade data. Surprisingly, despite these applications, the distributional properties of Hawkes type processes have not been thoroughly studied. We establish these properties below, and extend the Hawkes process to a broad family of self-exciting point processes.

We note two special cases. If $\kappa = 0$ in the specification (1), then λ is constant between events so J is a linear birth process. If $\delta = 0$, then J is a Poisson process whose intensity $u(t)$ evolves deterministically through time, and whose inter-arrival times are independent. Figure 3 contrasts the sample paths of the Poisson and Hawkes processes. It shows how the self-exciting property of the Hawkes process generates arrivals that are over-dispersed relative to Poisson arrivals. The resulting event clusters are strikingly similar to those of the empirical default process in Figure 1. The parameter δ in (1) allows us to directly control the frequency of these clusters. The parameter κ governs their magnitude.




Figure 2: Sample paths of the intensity (1) and the associated loss process L . A jump in the intensity represents the impact of a default. The jump size is the product of the loss at default and the sensitivity parameter $\delta = 1$. The loss at default is drawn from a uniform distribution on $\{0.4, 0.6, 0.8, 1\}$. The rate $\kappa = 5$ controls the exponential decay after an event. The reversion level $c = \lambda_0 = 0.7$.

While the Hawkes process induces clustered arrivals, it does not explode, and therefore the integral $A_t = \int_0^t \lambda_s ds$ is finite for all t . To see this, it suffices to consider the case $\kappa = 0$, when λ has non-decreasing sample paths. Then N is a linear birth process, which is non-explosive and integrable. From this we can conclude that N_t is integrable also in the case $\kappa > 0$. This, in turn, implies that A_t is integrable, and that the compensated local jump martingale $M = N - A$ is a martingale. To see this, we can appeal to Corollary 3 in Chapter II.6 of Protter (2004), which implies that M is a martingale if its quadratic variation is integrable. But the quadratic variation of M is equal to N . Further, we obtain that $E(M_t^2) = E(N_t) = E(A_t)$.

2.3 Dynkin formula

While the process J itself does not have the Markov property, the process $Y = (\lambda, J)^\top$ is a Markov process in a state space $D = \mathbb{R}_+ \times (\mathbb{R}_+ \times \mathbb{N})$. Consider the operator \mathcal{D} , acting on a function $g : D \rightarrow \mathbb{R}$ with continuous partial derivative $g_\lambda(\lambda, x)$, given by

$$(\mathcal{D}g)(\lambda, x) = \kappa(c - \lambda)g_\lambda(\lambda, x) + \lambda \int [g(\lambda + \delta z, x + (z, 1)^\top) - g(\lambda, x)] d\nu(z). \quad (3)$$

We provide an integrability condition guaranteeing that \mathcal{D} is the infinitesimal generator of Y . The generator describes the time evolution of the process over an infinitesimal time period and leads to a useful formula for expectations of functions of Y .




Figure 3: Sample paths of the Poisson and Hawkes processes. A bar indicates the number of arrivals in a given year. While the Poisson arrivals are evenly distributed over time due to the order statistics property, the Hawkes arrivals are clustered thanks to the self-exciting property. For the Hawkes process $c = \lambda_0 = 1$, $\kappa = 1.5$, $\delta = 2$ and the loss at default has a uniform distribution ν on $\{0.4, 0.6, 0.8, 1\}$. The Poisson process is the special Hawkes process for which $\delta = 0$ so that $\lambda_t = u(t)$. We choose $c = \lambda_0 = 10.57$ to match the expected number of Hawkes events over 30 years.

Proposition 2.1. Let $g : D \rightarrow \mathbb{R}$ be a function with continuous partial derivative $g_\lambda(\lambda, x)$ such that the variable

$$\int_0^t |(\mathcal{D}g)(\lambda_s, J_s) - \kappa(c - \lambda_s)g_\lambda(\lambda_s, J_s)| ds \quad (4)$$

is integrable for each t . Then the operator \mathcal{D} is the infinitesimal generator of the Markov process $Y = (\lambda, J)^\top$, and for $t \leq T$ we have a Dynkin formula

$$E(g(\lambda_T, J_T) | \mathcal{F}_t) = g(\lambda_t, J_t) + E\left(\int_t^T (\mathcal{D}g)(\lambda_s, J_s) ds \mid \mathcal{F}_t\right). \quad (5)$$

Proposition 2.1 provides a formula expressing the conditional expectation of a general function g of the Markov process $Y = (\lambda, J)^\top$ in terms of the infinitesimal generator (3). This formula is useful for functions g for which the expectation of $(\mathcal{D}g)(\lambda_s, J_s)$ can be calculated. We exploit it to obtain explicit formulas for the first and second moments of the intensity. Consider the function $g(\lambda, x) = \lambda$, which satisfies the integrability condition (4) of Proposition 2.1. We calculate

$$\begin{aligned} E((\mathcal{D}g)(\lambda_s, J_s)) &= E(\kappa(c - \lambda_s)) + E\left(\lambda_s \int \delta z d\nu(z)\right) \\ &= \kappa c + \mu E(\lambda_s) \end{aligned}$$

where $\mu = \delta \ell - \kappa$ and $\ell = \int z d\nu(z)$ is the expected loss at default. Letting $m(t) = E(\lambda_t)$, equation (5) implies that

$$m(t) = E(g(\lambda_t, J_t)) = \lambda_0 + \kappa c t + \mu \int_0^t m(s) ds. \quad (6)$$

In other words, the mean intensity solves the ordinary differential equation $m'(t) = \kappa c + \mu m(t)$ with initial condition $m(0) = \lambda_0$. The solution is

$$m(t) = E(\lambda_t) = \left(\frac{\kappa c}{\mu} + \lambda_0 \right) e^{\mu t} - \frac{\kappa c}{\mu} \quad (7)$$

provided that $\mu \neq 0$. Equation (7) shows that for $\mu > 0$, the mean intensity grows exponentially (without bound) in time. For $\mu \rightarrow 0$, the mean intensity grows linearly with time as $\lim_{\mu \rightarrow 0} E(\lambda_t) = \lambda_0 + \kappa c t$. For $\mu < 0$, the mean intensity decays exponentially with time to the long-run mean

$$\lim_{t \rightarrow \infty} E(\lambda_t) = -\frac{\kappa c}{\mu} > 0. \quad (8)$$

We apply a similar argument to the function $g(\lambda, x) = \lambda^2$ to show that the second moment $v(t) = E(\lambda_t^2)$ solves the ordinary differential equation

$$v'(t) = \rho m(t) + 2\mu v(t) \quad (9)$$

with initial condition $v(0) = \lambda_0^2$, where $\rho = 2\kappa c + \delta^2 \int z^2 d\nu(z)$. Here, we assume that ν has finite second moment. For $\mu \neq 0$, the solution to equation (9) is given by

$$v(t) = \frac{1}{2\mu^2} [\kappa c \rho (e^{\mu t} - 1)^2 + 2\mu \lambda_0 e^{\mu t} (\rho (e^{\mu t} - 1) + \mu \lambda_0 e^{\mu t})]. \quad (10)$$

We conclude that for $\mu < 0$, the long run variance of the intensity satisfies

$$\lim_{t \rightarrow \infty} \text{Var}(\lambda_t) = \lim_{t \rightarrow \infty} (v(t) - m^2(t)) = \frac{\kappa c \delta^2}{2\mu^2} \int z^2 d\nu(z). \quad (11)$$

Proposition 2.1 can be applied to the function $g(\lambda, x) = v \cdot (x_1^n, x_2^n)^\top$ to find a formula for the n th moment of J_t . For example, $E(v \cdot J_t) = v \cdot (\ell, 1)^\top \int_0^t m(s) ds$ with $m(t)$ given by equation (6), a formula that is foreshadowed by the more fundamental local martingale property of the compensated jump process $N - \int_0^\cdot \lambda_s ds$, which implies that $E(N_t) = \int_0^t E(\lambda_s) ds$. This argument does however not immediately lead to an explicit formula for higher order moments of J_t . Below, we calculate a transform of J , which has a wide range of applications, and which can be differentiated to obtain the moments of J .

2.4 Transform

Along with the distribution ν of the loss at default, the intensity specification (1) determines the distribution of the processes N and L . We develop a formula for a transform of the point process $J = (L, N)^\top$ that includes as special cases the Laplace and Fourier transforms. The transform can be inverted to obtain the corresponding distribution.

Let $u \in \mathbb{C}_-^2$, the set of tuples of complex numbers with non-positive real part. Express the conditional transform $E(\exp(u \cdot J_T) | \mathcal{F}_t)$ as $f(t, \lambda_t, J_t)$ for some complex-valued

function f on $[0, T] \times D$. For $f(t, \lambda_t, J_t)$ to be a martingale, its drift must vanish. This means that f must satisfy the partial integral differential equation

$$0 = f_t(t, \lambda, x) + (\mathcal{D}f)(t, \lambda, x)$$

with boundary condition $f(T, \lambda, x) = \exp(u \cdot x)$, where $(\mathcal{D}f)(t, \lambda, x)$ is obtained by applying the generator \mathcal{D} to the real and imaginary parts of $f(t, \cdot, \cdot)$. This PIDE reduces to a system of ordinary differential equations when f is taken to be an exponentially affine function of λ and x . This argument is made precise in the next result.

Proposition 2.2. *The transform of the point process $J = (L, N)^\top$ is given by*

$$E(\exp(u \cdot J_T) | \mathcal{F}_t) = \exp(a(t) + b(t)\lambda_t + u \cdot J_t) \quad (12)$$

where $t \leq T$, $u \in \mathbb{C}_-^2$ and the coefficient functions $a(t) = a(u, t, T)$ and $b(t) = b(u, t, T)$ satisfy the following ordinary differential equations:

$$\partial_t b(t) = \kappa b(t) + 1 - \theta(\delta b(t) + u \cdot (1, 0)^\top) \exp(u \cdot (0, 1)^\top) \quad (13)$$

$$\partial_t a(t) = -\kappa c b(t) \quad (14)$$

with boundary conditions $a(T) = b(T) = 0$, where θ is the jump transform

$$\theta(c) = \int e^{cz} d\nu(z), \quad c \in \mathbb{C}. \quad (15)$$

Proposition 2.2 states that the conditional transform of J_T given \mathcal{F}_t is an exponentially affine function of the intensity λ_t and the state J_t , with coefficient functions that satisfy a system of ODEs. To put this result into perspective and to prepare its extension, note that the intensity λ specified by (1) satisfies the equation

$$d\lambda_t = \kappa(c - \lambda_t)dt + \delta dL_t. \quad (16)$$

Equation (16) indicates that the drift of the Markov process $Y = (\lambda, J)^\top$ has affine dependence on Y . The jumps of Y arrive with an intensity that also has affine dependence on Y . It follows that Y is an affine jump diffusion process which has no diffusion term. But the transform of an affine jump diffusion is an exponentially affine function of its initial value, as is established by Duffie et al. (2000).

The argument behind Proposition 2.2 immediately extends to more general functionals of $Y = (\lambda, J)^\top$. For example, the argument leads to the transform of the vector Y , not just its component J . Section 4 covers this extension in a more general setting with multiple stochastic risk factors driving the intensity λ of J . We then obtain the joint transform of the point process and the risk factors. This yields, in particular, the transform of the intensity of the point process.

In some cases the ODEs (13) and (14) can be solved analytically, for example when $\kappa = 0$ and J is a birth process. In the general case, the ODEs are quickly solved numerically, using the Runge-Kutta algorithm for example.

2.5 Moments

By differentiating the conditional transform of J with respect to u and evaluating the derivative at $u = 0$, we find the conditional expectation

$$E(v \cdot J_T | \mathcal{F}_t) = \mathcal{A}(0, t, T) + \mathcal{B}(0, t, T)\lambda_t + v \cdot J_t \quad (17)$$

for $v \in \mathbb{R}^2$ and $t \leq T$, where the functions $\mathcal{A}(t) = \mathcal{A}(u, t, T)$ and $\mathcal{B}(t) = \mathcal{B}(u, t, T)$ satisfy

$$\partial_t \mathcal{B}(t) = \kappa \mathcal{B}(t) - (\delta \mathcal{B}(t) + v \cdot (1, 0)^\top) \theta'(\delta b(t)) - v \cdot (0, 1)^\top \theta(\delta b(t)) \quad (18)$$

$$\partial_t \mathcal{A}(t) = -\kappa c \mathcal{B}(t) \quad (19)$$

with boundary conditions $\mathcal{A}(T) = \mathcal{B}(T) = 0$. The derivative of the jump transform is

$$\theta'(c) = \int z e^{cz} d\nu(z), \quad c \in \mathbb{C},$$

and $a(t)$ and $b(t)$ satisfy the equations (13) and (14) and the relevant boundary conditions. Since $u = 0$, we can choose $a(t) = b(t) = 0$. Under mild assumptions on the distribution ν (continuity of $\theta'(c)$ suffices), this solution is unique and equation (18) simplifies to

$$\partial_t \mathcal{B}(t) = (\kappa - \delta \ell) \mathcal{B}(t) - v \cdot (\ell, 1)^\top \quad (20)$$

where $\ell = \int z d\nu(z)$ is the expected loss at default. Provided that $\mu = \delta \ell - \kappa \neq 0$, we obtain the following explicit solutions:

$$\mathcal{B}(t) = \frac{1}{\mu} v \cdot (\ell, 1)^\top (e^{\mu(T-t)} - 1) \quad (21)$$

$$\mathcal{A}(t) = \frac{1}{\mu} v \cdot (\ell, 1)^\top \kappa c \left(\frac{1}{\mu} (e^{\mu(T-t)} - 1) - T + t \right). \quad (22)$$

For $\mu \neq 0$, the mean number of events takes the form

$$E(N_t) = c_1 (e^{\mu t} - 1) + c_2 t$$

where $c_1 = (\kappa c + \mu \lambda_0) / \mu^2$ and $c_2 = -\kappa c / \mu$. If $\kappa = 0$ and N is a birth process, then $E(N_t)$ grows exponentially in t at rate $\delta \ell > 0$. If $\kappa > 0$, then the growth of $E(N_t)$ is determined by the sign of μ . If $\mu > 0$, then we have exponential growth at rate μ that is counteracted by linear decay at rate c_2 . If $\mu < 0$, then we have linear growth at rate $c_2 = \lim_{t \rightarrow \infty} E(\lambda_t)$ that is counteracted by exponential decay at rate μ .

Higher order conditional moments of J_T can be calculated similarly, by successively differentiating the transform. Thus, we obtain a closed form expression for conditional expectations of the form $E(f(J_T) | \mathcal{F}_t)$, where $f : \mathbb{R}_+ \times \mathbb{N} \rightarrow \mathbb{R}$ is an integrable function that is polynomial. We can estimate the conditional expectation $E(h(J_T) | \mathcal{F}_t)$ for any continuous, integrable function h on $\mathbb{R}_+ \times \mathbb{N}$ by approximating h with f .

2.6 Transform inversion

The transform of J_T can be inverted using Fast Fourier Transform techniques, to obtain the conditional distribution given \mathcal{F}_t of N_T and L_T for all future dates T . The distribution of N_T can also be obtained directly. Proposition 2.2 implies that the conditional probability generating function of the Hawkes process takes the form

$$E(v^{N_T - N_t} | \mathcal{F}_t) = \exp(c(t) + d(t)\lambda_t) \quad (23)$$

for $v \in (0, 1)$ and $t \leq T$, where the coefficient functions

$$c(t) = c(v, t, T) = a((0, \log v)^\top, t, T)$$

$$d(t) = d(v, t, T) = b((0, \log v)^\top, t, T)$$

satisfy the ordinary differential equations

$$\partial_t d(t) = \kappa d(t) + 1 - v\theta(\delta d(t)) \quad (24)$$

$$\partial_t c(t) = -\kappa c(t) \quad (25)$$

with boundary conditions $c(T) = d(T) = 0$. Equations (24)–(25) determine the distribution of the Hawkes process. Expansion of the left hand side of equation (23) into a power series shows that for $n = 0, 1, 2, \dots$

$$\varphi_t(n, T) = P(N_T - N_t = n | \mathcal{F}_t) = \frac{1}{n!} \partial_v^n \exp(c(v, t, T) + d(v, t, T)\lambda_t)|_{v=0} \quad (26)$$

In practice, the calculation of these probabilities calls for the solution of a system of ODEs derived from equations (24)–(25). In case $n = 0$ we get

$$\varphi_t(0, T) = \exp(C(t) + D(t)\lambda_t)$$

where the functions $D(t) = d(0, t, T)$ and $C(t) = c(0, t, T)$ satisfy the ODEs

$$\partial_t D(t) = \kappa D(t) + 1 \quad (27)$$

$$\partial_t C(t) = -\kappa C(t)$$

with boundary conditions $C(T) = D(T) = 0$. We obtain the formula

$$\varphi_t(0, T) = \exp\left(\frac{c - \lambda_t}{\kappa}(1 - e^{-\kappa(T-t)}) - c(T-t)\right) \quad (28)$$

provided that $\kappa > 0$. Note that the probability (28) of no events during $(t, T]$ does not depend on the clustering parameter δ . It agrees with the probability of no events during $(t, T]$ for a time-inhomogeneous Poisson process with intensity $u(t) = c + (\lambda_0 - c) \exp(-\kappa t)$. This is because between events, the Hawkes intensity (1) is deterministic and governed by the function $u(t)$.

For $n \geq 1$ the probability (26) depends on the clustering parameter δ . Using Faà di Bruno's formula, we get

$$\varphi_t(n, T) = \varphi_t(0, T) \sum \frac{1}{m_1! \dots m_n!} \prod_{k=1}^n \left(\frac{C_k(t) + D_k(t)\lambda_t}{k!} \right)^{m_k} \quad (29)$$

where the sum is over all n -tuples (m_1, \dots, m_n) of non-negative integers that satisfy the constraint $m_1 + 2m_2 + \dots + nm_n = n$ and the functions

$$D_k(t) = \partial_v^k d(v, t, T)|_{v=0}$$

$$C_k(t) = \partial_v^k c(v, t, T)|_{v=0}$$

satisfy the ordinary differential equations

$$\partial_t D_k(t) = \kappa D_k(t) - k E_{k-1}(t) \quad (30)$$

$$\partial_t C_k(t) = -\kappa C_k(t) \quad (31)$$

with boundary conditions $C_k(T) = D_k(T) = 0$, where $E_k(t) = \partial_v^k \theta(\delta d(v, t, T))|_{v=0}$ can be calculated by another application of Faà di Bruno's formula:

$$E_k(t) = \sum \frac{k!}{m_1! \dots m_k!} \theta^{(m_1 + \dots + m_k)}(\delta D(t)) \prod_{i=1}^k \left(\frac{\delta D_i(t)}{i!} \right)^{m_i}$$

and the sum is over all k -tuples (m_1, \dots, m_k) of non-negative integers that satisfy $m_1 + 2m_2 + \dots + km_k = k$. The equations (30) and (31) are solved numerically unless $\delta = 0$, in which case N is a time-inhomogeneous Poisson process with intensity $u(t)$.

3 Portfolio credit derivatives

We show how to use the results developed above to analyze portfolio credit derivatives, which expose an investor to the risk of correlated default in a reference portfolio of credit-sensitive instruments such as loans or bonds. We consider index and tranche swaps, which are the most popular portfolio derivatives. These are bilateral contracts, in which one party provides default protection on the reference portfolio, and the other party pays a premium for this protection. The basic valuation problem, addressed below, is to determine the fair premium, which is governed by the correlated default timing in the reference portfolio. Through a market calibration, we show that the basic intensity model (1) captures the correlated default risk implied by the premia offered in the market.

3.1 Valuation

Index and tranche swaps are based on a portfolio whose n constituent securities have a notional 1, maturity date T and premium payment dates (t_m) . The loss at default

$\ell_n \in [0, 1]$. The swap is specified by a lower attachment point $\underline{K} \in [0, 1]$ and an upper attachment point $\overline{K} \in (\underline{K}, 1]$. An index swap has attachment points $\underline{K} = 0$ and $\overline{K} = 1$. The swap notional $K = n(\overline{K} - \underline{K})$. The protection seller covers portfolio losses as they occur, given that the cumulative losses are larger than \underline{K} but do not exceed \overline{K} . The cumulative payments at time t , denoted U_t , are given by the “call spread”

$$U_t = (L_t - \underline{K}n)^+ - (L_t - \overline{K}n)^+. \quad (32)$$

The value at time $t \leq T$ of these payments is given by

$$D_t = E \left(\int_t^T e^{-r(s-t)} dU_s \mid \mathcal{F}_t \right) \quad (33)$$

where, here and below, the reference measure P is a risk-neutral pricing measure with respect to an interest rate $r > 0$. By Stieltjes integration by parts, we can conveniently express the value D_t in terms of conditional expectations of U :

$$D_t = e^{-r(T-t)} E(U_T \mid \mathcal{F}_t) - U_t + r \int_t^T e^{-r(s-t)} E(U_s \mid \mathcal{F}_t) ds. \quad (34)$$

The protection buyer receives the loss payments, and in return, makes premium payments to the protection seller. Each premium payment has two parts. The first part is an upfront payment, which is expressed as a fraction F of the tranche notional K . For an index swap, $F = 0$. The second part consists of payments that are proportional to the premium notional I_t , which is given by $n - (N_t \wedge n) = n + (N_t - n)^+ - N_t$ for an index swap and $K - U_t$ for a tranche swap with $\overline{K} < 1$. Let c_m be the day count fraction for the period m , roughly 1/4 for quarterly payments. Then, with S denoting the running premium rate, the value at time $t \leq T$ of the premium payments is given by

$$P_t(F, S) = FK + S \sum_{t_m \geq t} e^{-r(t_m-t)} c_m E(I_{t_m} \mid \mathcal{F}_t). \quad (35)$$

For a fixed upfront rate F , the running spread S_t at time t is the solution $S = S_t$ to the equation $D_t = P_t(F, S)$. Setting $D_t = P_t(F, S)$ for a fixed S gives a value $F = F_t$ for the time t upfront rate F_t . Formulae (34) and (35) indicate that these rates depend only on call options $E((v \cdot J_s - c)^+ \mid \mathcal{F}_t)$ with various strikes c , maturities $s \in (t, T]$, and values $v \in \{(0, 1)^\top, (1, 0)^\top\}$. Thus, to value a swap we need only calculate the values of options on N and L . One approach to calculate these values is to integrate the option payoff function $(x - c)^+$ against the point process distribution. The latter is obtained by inverting the point process transform in Proposition 2.2. In the case of N , the inverse transform is given by formula (26). Alternatively, we can invert the transform of the option price, which is a function of the transform of the point process and that of the payoff function, see Lee (2004). We may also apply the saddlepoint approximations developed by Glasserman & Kim (2008) for affine jump diffusion models. These are applicable here since the transform of J_s is an exponentially affine function of the intensity.




Figure 4: Annualized premium for protection against default losses in a portfolio of 100 equally weighted securities as implied by the model specification (1), stated in basis points (1 basis point = 10^{-4}). We assume quarterly premium payments, and set $\lambda_0 = c = 1$. The risk-free rate $r = 5\%$. The loss at default ℓ_n is uniformly distributed on $\{0.24, 0.96\}$ with expectation $\ell = 0.6$. *Left panel*: Index swap spread as a function of the maturity date T , for each of several values of the clustering parameter δ . The decay rate $\kappa = 1$. *Right panel*: 5 year index spread as a function of δ , for each of several values of κ .

If the distribution of J_s is light-tailed, then for the relatively large n in practice,¹ the distribution of $(v \cdot J_s) \wedge n = v \cdot J_s - (v \cdot J_s - n)^+$ may be well approximated by the distribution of $v \cdot J_s$ for $v \in \{(0, 1)^\top, (1, 0)^\top\}$. In this case, the value $E((v \cdot J_s - n)^+ | \mathcal{F}_t)$ is negligible, and the index swap spread depends only on $E(v \cdot J_s | \mathcal{F}_t)$. Thus, thanks to formula (17), we obtain an approximate closed formula for the index swap spread.

3.2 Numerical examples

We provide numerical examples of index and tranche rates to develop some intuition for the model parameters. Based on the approximate index swap spread formula, the left panel of Figure 4 shows the index swap spread S_0 as a function of the maturity date T , for each of several values of the clustering parameter δ . For $\delta = 0$, the portfolio loss process is a compound Poisson process, which generates a flat spread term structure. All else fixed, the higher δ , the more frequent the event clusters, the higher the spread and the steeper the term structure. The right panel of Figure 4 shows the $T = 5$ year index swap spread as a function of δ , for each of several values of the decay rate κ . For $\kappa = 0$, the portfolio loss process is a compound birth process. All else equal, the higher κ , the faster the impact of an event decays, and the smaller the chance of large losses.

The tranche swap rate increases with maturity and decreases with seniority. An

¹The standard portfolio size is 100 or 125. Bespoke deals can reference substantially larger portfolios.




Figure 5: Upfront premium rate for protection against default losses in a tranche of a portfolio of 100 equally weighted securities as implied by the model specification (1). We assume the running spread is equal to zero. The maturity $T = 5$ years. We set $\lambda_0 = c = 1$ and $\kappa = 1$. The risk-free rate $r = 5\%$. *Left panel:* Tranche upfront rate as a function of the clustering parameter δ , for each of several sets of standard attachment points. The loss at default ℓ_n is uniformly distributed on $\{0.24, 0.96\}$ with expectation $\ell = 0.6$. *Right panel:* Tranche upfront rate as a function of the volatility of the loss at default ℓ_n , for each of several sets of standard attachment points. The expected loss at default is 0.6. The clustering parameter $\delta = 1$.

increase in the lower attachment point, \underline{K} , leads to greater subordination (that is, buffer capital) available to the protection seller, who covers only the losses in excess of \underline{K} . The left panel of Figure 5 shows the upfront tranche swap rate² F_0 as a function of the clustering parameter δ for each of several sets of standard attachment points, assuming the running spread $S_0 = 0$. The sensitivity of the tranche rate to δ increases with the seniority of the tranche. This is because senior tranches are most exposed to default clustering, the frequency of which is controlled by δ . The bigger δ , the fatter the tail of the portfolio loss distribution. Since relatively few defaults suffice to wipe out the subordinated tranches, the frequency of clusters is less relevant for the pricing of these tranches. The right panel of Figure 5 shows the upfront tranche rate as a function of the volatility of the loss at default ℓ_n . The sensitivity to the volatility of the loss increases with seniority.

3.3 Market calibration

We perform market calibration experiments to illustrate the empirical importance of the features of our basic intensity model. To this end, we fit the parameters of the specification (1) from index and tranche swap market rates for each of the 21 trading days of September

²See Section 3.3 for details on the method of computation.

2008. This month witnessed significant volatility due to the demise of Fannie Mae and Freddy Mac on the 8th, the default of Lehman Brothers on the 15th, the collapse of American International Group on the 16th, the problems appearing at Morgan Stanley on the 18th, and the default of Washington Mutual on the 25th. The swap market rates are for the $T = 5$ year maturity, and are obtained from UBS. They reference the CDX High Yield portfolio, which consists of 100 equally weighted names of relatively low credit quality. The tranches have attachment points $(0, 10\%)$, $(10\%, 15\%)$, $(15\%, 25\%)$ and $(25\%, 35\%)$. Thus, together with the quote for the index contract, we have 5 quotes per calibration date, from which we fit the parameter vector $\theta = (c, \kappa, \delta, \lambda_0)$. In accordance with market practice, we assume that the distribution of the loss at default $\nu = \delta_{0.6}$. The risk-free rate of interest r is set to 5%. Swap premium payments are made quarterly. Using a gradient-based method, we numerically solve the nonlinear optimization problem

$$\min_{\theta \in \Theta} \sum_m \frac{(\text{Mid}(m) - \text{Model}(m, \theta))^2}{\text{Mid}(m)} \quad (36)$$

where $\Theta = (0, 8] \times [0, 10] \times [0, 20] \times (0, 30]$ and the sum ranges over the contracts. The market mid-quote $\text{Mid}(m)$ is the arithmetic average of the market bid and ask quotes for index or tranche m . The model rate for index or tranche m is denoted $\text{Model}(m, \theta)$, and is given by the formulae developed in Section 3.1. The optimization is initialized at a set of parameter values drawn from a uniform distribution over the parameter space Θ , and is repeated for each of 100 independent draws. The optimal parameter vector θ^* is the solution to (36) with the minimum objective function value among all 100 runs.

The index and tranche swap pricer and the optimization are implemented in Matlab. The ODEs (13)–(14) determining the point process transform (12) are solved numerically using the Runge-Kutta method. The transform (12) is inverted numerically using the Fast Fourier Transform. We use 1100 sample points, a value we found to be a good tradeoff between speed and accuracy. To get a sense of the numerical accuracy of this procedure, we contrast the distribution of N generated by the FFT with that based on formula (29). We find that the FFT produces very accurate results. It is also quicker than the evaluation of formula (26), which requires high-precision algorithms. The computations are performed on a PC with a 2.66 GHz Intel Processor and 4 GB of RAM.

For each of the 21 calibration dates, we calculate the average absolute percentage pricing error (AAPE), given by $(1/5) \sum_{m=1}^5 |\text{Model}(m, \theta^*) - \text{Mid}(m)| / \text{Mid}(m)$. The left panel of Figure 6 shows the time series of these errors during September 2008. The model fits the market on each date, with an acceptable average AAPE over all calibration dates of 4.4%, and a variance of 0.086. For comparison, the market standard copula model completely failed calibration on several days during September 2008.³ The calibration errors are relatively low during the most volatile period, which started on September 15th with the default of Lehman. This indicates that the model captures the default correlation

³That is, calibration required parameter values outside the set of admissible model parameter values. The copula model called for a correlation coefficient larger than 1, which is not meaningful.




Figure 6: Calibration results for the basic intensity model (1) for the 21 trading days of September 2008, for the CDX High Yield portfolio. *Left Panel:* Average absolute percentage error (AAPE) for each calibration date. *Right Panel:* Value of the clustering parameter δ for each calibration date.

implied by the market prices, especially in periods of extreme stress. The calibrated values of the clustering parameter δ , shown in the right panel of Figure 6, suggest that this is related to the self-exciting property of the default process. The time series behavior of the calibrated values of δ clearly reflects the events in mid September. It indicates the fear of investors of a cluster of events triggered by the default of Lehman on the 15th (first peak) and the near collapse of Morgan Stanley on the 18th (second peak).

4 Affine point processes

We extend the basic model specification analyzed in the preceding two sections. Our primary objective is to permit a richer structure for the intensity without reducing the computational tractability of the basic specification. The extension is required for applications in which the intensity is influenced by a set of risk factors that follow stochastic processes on their own. In the basic model (1), the point process itself is the only such risk factor. The extension facilitates the inclusion of exogenous (jump-) diffusion risk factors that are relevant to event arrivals. This is particularly important for empirical applications, in which the intensity model is estimated from a time series of portfolio derivative market prices, as in Azizpour & Giesecke (2008) and Longstaff & Rajan (2008). In these applications, the model must replicate the diffusive fluctuation of market prices, and this requires the presence of diffusive risk factors.

The idea behind the extension is to replace the intensity component λ of the Markov process $(\lambda, J)^\top$ analyzed in Proposition 2.1 by a Markov process X that represents a

vector of stochastic risk factors. The process X drives the intensity of the jumps of J . The transform of $(X, J)^\top$ is computationally tractable if X is taken to be an affine jump diffusion. This formulation leads to an affine point process.

4.1 Specification

We call a point process *affine* if its event arrival intensity is an affine function of an affine jump diffusion and its jump sizes are drawn from a fixed distribution. A Markov process X in a state space $D \subset \mathbb{R}^d \times \mathbb{R}_+$ is an affine jump diffusion in the sense of Duffie et al. (2000) if X is a strong solution to the stochastic differential equation

$$dX_t = \mu(X_t, t) dt + \sigma(X_t, t) dW_t + \sum_{i=1}^m \zeta^i dZ_t^i, \quad X_0 \in \mathbb{R}^d \quad (37)$$

where W is an \mathbb{R}^d -valued standard Brownian motion, $\mu : D \rightarrow \mathbb{R}^d$ is the drift, $\sigma : D \rightarrow \mathbb{R}^{d \times d}$ is the volatility and each Z^i is a *temporally consistent* \mathbb{R}_+^d -valued point process. In other words, the component processes of each vector Z^i share event times and differ only in jump sizes, and we denote their common intensity by $\lambda^i(X_t, t)$ for some $\lambda^i : D \rightarrow \mathbb{R}_+$. The jump sizes are drawn from a distribution ν^i on \mathbb{R}_+^d that has no mass zero at 0. Each parameter ζ^i is a d -dimensional diagonal matrix. We assume that for each t , $\{x : (x, t) \in D\}$ contains an open subset of \mathbb{R}^d . For time-dependent coefficient functions that are bounded and continuous on \mathbb{R}_+ , we assume that

$$\begin{aligned} \mu(x, t) &= K_0(t) + K_1(t)x, \quad K_0(t) \in \mathbb{R}^d, \quad K_1(t) \in \mathbb{R}^{d \times d} \\ (\sigma(x, t)\sigma(x, t)^\top)_{jk} &= (H_0)_{jk}(t) + (H_1)_{jk}(t) \cdot x, \quad H_0(t) \in \mathbb{R}^{d \times d}, \quad H_1(t) \in \mathbb{R}^{d \times d \times d} \\ \lambda^i(x, t) &= \Lambda_0^i(t) + \Lambda_1^i(t) \cdot x, \quad \Lambda_0^i(t) \in \mathbb{R}, \quad \Lambda_1^i(t) \in \mathbb{R}^d, \quad i = 1, 2, \dots, m. \end{aligned}$$

Whether an affine point process J driven by X has the self-exciting property depends on the relation between J and its intensity $\lambda_t = \lambda(X_t, t)$, where $\lambda : D \rightarrow \mathbb{R}_+$. The self-exciting property holds if X , and hence λ , depends on J itself. A sufficient condition is that at least one of the component processes of J is temporally consistent with one of the component processes of one of the jump terms Z^i of X . Since the intensity is a function of the realized loss at default, we also generate a dependence structure between default and recovery rates in the self-exciting case.

4.2 Self-exciting examples

To illustrate the specification of an affine point process, we first consider the two dimensional case $J = (L, N)^\top$. J is driven by a one dimensional risk factor X with a single jump term Z that we identify with the component process L . The remaining component process N is temporally consistent with Z . Thus, L and N are one dimensional affine point processes that share common event times that arrive with intensity $\lambda(X_t, t)$. By

construction, the jump sizes of L are governed by the distribution ν . The jumps of N are unit-sized since N counts the arrivals. The specification of J as an affine point process is completed with the specification of the coefficient functions of the risk factor X .

Example 4.1. *Suppose $K_0(t) = \kappa c$ for $\kappa \geq 0$ and $c > 0$, $K_1(t) = -\kappa$, $H_0(t)$ is a matrix of zeros, $H_1(t)$ is a tensor of zeros, $X_0 = c$ and $\zeta = \delta \geq 0$. Let $\Lambda_0(t) = 0$ and $\Lambda_1(t) = 1$. Then the intensity $\lambda = X$ of J satisfies the basic model (1):*

$$d\lambda_t = \kappa(c - \lambda_t) dt + \delta dL_t.$$

A significant generalization of the basic specification is made by introducing Brownian terms and independent jump terms in the intensity. The Brownian terms model the diffusive fluctuation in the default rate and can be driven by a stochastic volatility. The jump terms model the sensitivity of the intensity to market events, such as macro-economic shocks or defaults to names that are outside the portfolio.

Example 4.2. *Suppose the coefficient functions of X are chosen as in Example 4.1, with the exception of the tensor $H_1(t)$, for which $(H_1)_{111}(t) = \sigma^2$ for $\sigma \geq 0$; all other elements are zero. Then the intensity $\lambda = X$ of J satisfies the stochastic differential equation*

$$d\lambda_t = \kappa(c - \lambda_t) dt + \sigma \sqrt{\lambda_t} dW_t + \delta dL_t, \quad (38)$$

showing that between events, the intensity drifts stochastically toward c with diffusive fluctuations driven by W . A sample path of the intensity is in Figure 7. Compare with the sample path of the basic model (1) in Figure 2. The extended model (38) is estimated from time series of index and tranche swap rates by Azizpour & Giesecke (2008).

In practice, the impact of a default on the other firms may depend on the characteristics of the defaulter. The next example allows us to distinguish between different types of firms represented in the portfolio. We consider a two dimensional affine point process $J = (L^1, L^2)^\top$ whose components record portfolio losses triggered by defaults of two firm types. The portfolio loss process L is the sum $L^1 + L^2$. The process J is driven by a two dimensional risk factor $X = (X^1, X^2)^\top = (\lambda^1, \lambda^2)^\top$ with two jump terms Z^1 and Z^2 whose components are identified respectively with L^1 and L^2 . This means, that both component processes of Z^i are indistinguishable from L^i .

Example 4.3. *Suppose*

$$K_0(t) = \begin{pmatrix} \kappa^1 c^1 \\ \kappa^2 c^2 \end{pmatrix}, \quad K_1(t) = \begin{pmatrix} -\kappa^1 & 0 \\ 0 & -\kappa^2 \end{pmatrix},$$

$H_0(t)$ is a matrix of zeros, $H_1(t)$ is a tensor of zeros and

$$\zeta^1 = \begin{pmatrix} \delta^{1,1} & 0 \\ 0 & \delta^{2,1} \end{pmatrix}, \quad \zeta^2 = \begin{pmatrix} \delta^{1,2} & 0 \\ 0 & \delta^{2,2} \end{pmatrix}.$$




Figure 7: Sample paths of the intensity (38) and the associated loss process L . A jump in the intensity represents the impact of a default. The jump size is the product of the loss at default and the sensitivity parameter $\delta = 1$. The loss at default is drawn from an independent uniform distribution on $\{0.4, 0.6, 0.8, 1\}$. The reversion rate $\kappa = 5$ and the reversion level $c = \lambda_0 = 0.7$. The volatility $\sigma = 0.2$ controls the diffusive fluctuation of the intensity between events.

Let $\Lambda_0^i(t) = 0$ for $i = 1, 2$, $\Lambda_1^1(t) = (1, 0)^\top$ and $\Lambda_1^2(t) = (0, 1)^\top$. Then the intensities $\lambda^1 = X^1$ and $\lambda^2 = X^2$ satisfy the stochastic differential equations

$$\begin{aligned} d\lambda_t^1 &= \kappa^1(c^1 - \lambda_t^1) dt + \delta^{1,1} dL_t^1 + \delta^{1,2} dL_t^2 \\ d\lambda_t^2 &= \kappa^2(c^2 - \lambda_t^2) dt + \delta^{2,1} dL_t^1 + \delta^{2,2} dL_t^2. \end{aligned}$$

While the parameters $\delta^{1,1}$ and $\delta^{2,2}$ control the self-excitation, the parameters $\delta^{1,2}$ and $\delta^{2,1}$ control the cross-excitation of the processes L^1 and L^2 .

The preceding example illustrates the significance of allowing J to be vector-valued. The multi dimensional setting facilitates the specification of (univariate) point processes whose arrivals are correlated. The dependence between the components of J can be induced by a diffusion risk factor that is common to the component intensities. It can also be generated by direct interaction terms, as in Example 4.3. In the latter case, an arrival of one component process has an impact on the intensities of other component processes. This facilitates the modeling of cross-excitation phenomena. For example, the component processes can model distinct portfolios. A default causes a loss in the respective portfolio, and also has an impact on firms in the other portfolios.

4.3 Transform

We show that a general affine point process J is as computationally tractable as the basic specification (1). To derive a formula for the transform of J , we can apply the PIDE arguments developed in Propositions 2.1 and 2.2, which are based on the generator of the Markov process $Y = (X, J)^\top$. We propose an alternative approach that consists of constructing Y as an affine jump diffusion process in an enlarged state space, and then applying to this process the transform characterization for affine jump diffusions developed by Duffie et al. (2000). Our motivation for pursuing this approach is twofold. First, we wish to provide an alternative to our earlier PIDE arguments. Second, we would like to integrate our point process model formulation into the affine jump diffusion setting, which is standard in many areas.

Proposition 4.4. *Suppose J is a dimension k affine point process driven by a dimension d affine jump diffusion X with state space $D^X \times \mathbb{R}_+ \subset \mathbb{R}^d \times \mathbb{R}_+$. Then $Y = (X, J)^\top$ is a dimension $(k + d)$ affine jump diffusion with state space $D^Y \times \mathbb{R}_+ \subset \mathbb{R}^{k+d} \times \mathbb{R}_+$ whose coefficients $(K_0, K_1, H_0, H_1, \Lambda_0^i, \Lambda_1^i, \zeta^i)$ and jump distributions ν^i are canonically determined by the corresponding items for X .*

In Proposition 4.4 and below, we use the same symbols for the coefficients of X and Y to avoid further complicating notation. We illustrate the idea on Example 4.3.

Example 4.5. *In the setting of Example 4.3, the four dimensional affine jump diffusion $Y = (X, J)^\top = (\lambda^1, \lambda^2, L^1, L^2)^\top$ has two (four dimensional) jump terms Z^1 and Z^2 whose components are indistinguishable from L^1 and L^2 respectively. Since the third and fourth components of Y are driftless, the third and fourth rows of the drift coefficients are populated with zeros. These coefficients are given by*

$$K_0(t) = (\kappa^1 c^1, \kappa^2 c^2, 0, 0)^\top, \quad K_1(t) = \text{diag}(-\kappa^1, -\kappa^2, 0, 0).$$

The volatility coefficients $H_0(t)$ and $H_1(t)$ are zero since X has no Brownian term. The sensitivity matrices are

$$\zeta^1 = \text{diag}(\delta^{1,1}, \delta^{2,1}, 1, 0), \quad \zeta^2 = \text{diag}(\delta^{1,2}, \delta^{2,2}, 0, 1).$$

Finally, we give coefficients of the arrival intensities $\lambda_t^i = \Lambda_0^i(t) + \Lambda_1^i(t) \cdot Y_t$. We have $\Lambda_0^1(t) = \Lambda_0^2(t) = 0$ and

$$\Lambda_1^1(t) = (1, 0, 0, 0)^\top, \quad \Lambda_1^2(t) = (0, 1, 0, 0)^\top.$$

Proposition 4.4 facilitates the application of Proposition 1 in Duffie et al. (2000), which expresses the conditional transform of an affine jump diffusion as an exponentially affine function of its current value. Let \mathbb{C}^n denote the set of n -tuples of complex numbers.

When well defined at $t \leq T$ and $u \in \mathbb{C}^{d+k}$, the conditional transform of the $(d+k)$ dimensional affine jump diffusion $Y = (X, J)^\top$ constructed in Proposition 4.4 is given by

$$\psi(u, Y_t, t, T) = E(\exp(u \cdot Y_T) | \mathcal{F}_t). \quad (39)$$

Under technical conditions stated in Proposition A.1 in the appendix,

$$\psi(u, Y_t, t, T) = \exp(\alpha(u, t, T) + \beta(u, t, T) \cdot Y_t) \quad (40)$$

where the coefficient functions $\beta(t) = \beta(u, t, T)$ and $\alpha(t) = \alpha(u, t, T)$ satisfy the ordinary differential equations

$$\partial_t \beta(t) = -K_1(t)^\top \beta(t) - \frac{1}{2} \beta(t)^\top H_1(t) \beta(t) - \sum_{i=1}^m \Lambda_1^i(t) (\theta^i(\zeta^i \beta(t)) - 1) \quad (41)$$

$$\partial_t \alpha(t) = -K_0(t) \cdot \beta(t) - \frac{1}{2} \beta(t)^\top H_0(t) \beta(t) - \sum_{i=1}^m \Lambda_0^i(t) (\theta^i(\zeta^i \beta(t)) - 1) \quad (42)$$

with boundary conditions $\alpha(T) = 0$ and $\beta(T) = u$ and jump transforms

$$\theta^i(c) = \int_{\mathbb{R}_+^{d+k}} e^{c \cdot z} d\nu^i(z). \quad (43)$$

Under additional technical conditions stated in Proposition A.2 in the appendix, the transform (40) can be differentiated with respect to u to get the formula

$$E(e^{u \cdot Y_T} v \cdot Y_T | \mathcal{F}_t) = \psi(u, Y_t, t, T) (A(u, v, t, T) + B(u, v, t, T) \cdot Y_t) \quad (44)$$

where $t \leq T$, $u \in \mathbb{C}^{d+k}$, and $v \in \mathbb{R}^{d+k}$, and where the coefficient functions $B(t) = B(u, v, t, T)$ and $A(t) = A(u, v, t, T)$ satisfy the ordinary differential equations

$$\partial_t B(t) = -K_1(t)^\top B(t) - \beta(t)^\top H_1(t) B(t) - \sum_{i=1}^m \Lambda_1^i(t) \nabla \theta^i(\zeta^i \beta(t)) \cdot \zeta^i B(t) \quad (45)$$

$$\partial_t A(t) = -K_0(t) \cdot B(t) - \beta(t)^\top H_0(t) B(t) - \sum_{i=1}^m \Lambda_0^i(t) \nabla \theta^i(\zeta^i \beta(t)) \cdot \zeta^i B(t) \quad (46)$$

with boundary conditions $A(T) = 0$ and $B(T) = v$. As shown in Corollary A.3 in the appendix, in case $u = 0$ these differential equations admit the solutions⁴

$$B(0, v, t, T) = v \exp \left(\int_t^T \left(K_1(s)^\top + \sum_{i=1}^m \Lambda_1^i(s) \otimes \ell^i \zeta^i \right) ds \right) \quad (47)$$

$$A(0, v, t, T) = \int_t^T \left(K_0(s) + \sum_{i=1}^m \Lambda_0^i(s) \ell^i \zeta^i \right) \cdot B(s) ds, \quad (48)$$

⁴The tensor product $u \otimes v$ of vectors u and v is a matrix whose ij th element is $u_i v_j$.

which lead to a closed formula for $E(v \cdot Y_T | \mathcal{F}_t)$. Under additional technical conditions, the quantity (44) can be differentiated further to obtain higher order moments of Y_T .

The transform (40) encodes the *joint* distribution of the risk factor process X and the point process J . It facilitates applications that require the calculation of functionals of the form $E(h(X_T, J_T) | \mathcal{F}_t)$ for suitable h . An example is the valuation of forward and option contracts on index and tranche swaps, see Ding et al. (2006).

We can follow the argument in Section 2.6 to characterize in terms of ODEs derived from equations (41)–(42), the probability distribution of an integer-valued affine point process that may be driven by a multi-dimensional risk factor process X . Alternatively, we can apply Fourier inversion to the transform, which would also cover multi dimensional and real-valued affine point processes.

4.4 Further extensions

The analysis in the previous sections can be extended along several dimensions. We mention extensions that do not require new results. The specification can be generalized to include a stochastic discount rate that is driven by the affine jump diffusion Y . Since Y includes the affine point process J , this specification would generate a dependence structure among default, recovery and risk-free rates. Another potential extension is the generalization of the driving risk factor process X to a more general Markov process, such as the general affine process analyzed by Duffie, Filipovic & Schachermayer (2003) or the linear-quadratic jump diffusion process analyzed by Cheng & Scaillet (2007).

5 Conclusion

We specify and analyze a family of multivariate point processes whose arrival intensity is driven by an affine jump diffusion process. The component processes are self- and cross-exciting, and allow the description of comprehensive feedback effects among arrivals of different types. While these affine point processes significantly generalize the Poisson, birth and Hawkes processes, they offer a high degree of computational tractability in applications. We show that ordinary differential equations characterize the transform of an affine point process and the probability distribution of an integer-valued affine point process. The moments of an affine point process take a closed form.

Affine point processes have applications in many important areas, including finance, insurance, reliability, health care, and queuing. We illustrate their application to the valuation of portfolio credit derivatives, which are securities with payoffs that depend on the cumulative loss due to default in a portfolio of corporate bonds or loans. Here, an affine point process describes the time evolution of the portfolio loss. This specification incorporates the correlated timing of default, which is due to the exposure of bond issuers to common risk factors, and the impact of a default on the surviving issuers. It also incorporates the negative correlation between default and recovery rates. Our transform results

lead to computationally tractable valuation relations for portfolio credit derivatives, which we illustrate for index and tranche swaps.

A Technical results and proofs

Proof of Proposition 2.1. Note that Y is a process with right continuous paths of finite variation. By change of variables for Stieltjes integrals,

$$g(\lambda_t, J_t) - g(\lambda_0, J_0) = \int_0^t g_\lambda(\lambda_s, J_s) \kappa(c - \lambda_s) ds + \sum_{0 < s \leq t} [g(\lambda_s, J_s) - g(\lambda_{s-}, J_{s-})].$$

We can write

$$\begin{aligned} \sum_{0 < s \leq t} [g(\lambda_s, J_s) - g(\lambda_{s-}, J_{s-})] &= \int_0^t [g(\lambda_s, J_s) - g(\lambda_{s-}, J_{s-})] dN_s \\ &= \int_0^t \int [g(\lambda_{s-} + \delta z, J_{s-} + (z, 1)^\top) - g(\lambda_{s-}, J_{s-})] d\nu(z) dN_s \\ &= U_t^g + \int_0^t \int [g(\lambda_s + \delta z, J_s + (z, 1)^\top) - g(\lambda_s, J_s)] d\nu(z) \lambda_s ds \end{aligned}$$

where U^g is a process defined by

$$U_t^g = \int_0^t \int [g(\lambda_{s-} + \delta z, J_{s-} + (z, 1)^\top) - g(\lambda_{s-}, J_{s-})] d\nu(z) dM_s$$

and $M = N - \int_0^\cdot \lambda_s ds$ is a martingale. The integrability condition on the predictable integrand guarantees that U^g is a martingale, see Theorem 8 in Chapter II of Brémaud (1981). We conclude that $g(\lambda, J)$ is a special semimartingale with unique decomposition into a sum of a predictable finite variation process and a martingale:

$$g(\lambda_t, J_t) = g(\lambda_0, J_0) + \int_0^t (\mathcal{D}g)(\lambda_s, J_s) ds + U_t^g.$$

Since U^g is a martingale, so is the process defined by

$$g(\lambda_t, J_t) - g(\lambda_0, J_0) - \int_0^t (\mathcal{D}g)(\lambda_s, J_s) ds,$$

and this yields formula (5). □

Proof of Proposition 2.2. For fixed T and $u \in \mathbb{C}_-^2$, define the function $f : [0, T] \times D \rightarrow \mathbb{C}$ by the conditional expectation

$$f(t, \lambda, x) = E(\exp(u \cdot J_T) \mid \lambda_t = \lambda, J_t = x).$$

Then $E(\exp(u \cdot J_T) | \mathcal{F}_t) = f(t, \lambda_t, J_t)$. Since for $u \in \mathbb{C}_-^2$ the real and imaginary parts of the function $g(\lambda, x) = \exp(u \cdot x)$ are in the domain of the generator \mathcal{D} so are the real and imaginary parts of the function $f(t, \lambda, x)$, for each time t . The function $f(t, \lambda, x)$ is also differentiable with respect to t , and the derivative is continuous, see Proposition VII.1.2 in Revuz & Yor (2005). For this, we note that $Y = (\lambda, J)^\top$ is a Feller process.

Now, for $(f(t, \lambda_t, J_t))_{t \leq T}$ to be a martingale, we must have that

$$\begin{cases} f_t(t, \lambda, x) + (\mathcal{D}f)(t, \lambda, x) = 0 \\ f(T, \lambda, x) = \exp(u \cdot x) \end{cases} \quad (49)$$

where $(\mathcal{D}f)(t, \lambda, x)$ is obtained by applying \mathcal{D} to the real and imaginary parts of $f(t, \cdot, \cdot)$. We propose the following change of variables:

$$w(t, \lambda, u) = f(t, \lambda, x) e^{-u \cdot x}$$

Then, after re-arranging terms and denoting $\bar{z} = (z, 1)^\top$, equation (49) takes the form

$$\begin{cases} w_t(t, \lambda, u) + \lambda \int [w(t, \lambda + \delta z, u) - w(t, \lambda, u)] e^{u \cdot \bar{z}} d\nu(z) \\ + \lambda \int (e^{u \cdot \bar{z}} - 1) w(t, \lambda, u) d\nu(z) + \kappa(c - \lambda) w_\lambda(t, \lambda, u) = 0 \\ w(T, \lambda, u) = 1 \end{cases}$$

This PIDE has an affine form in λ . We take $w(t, \lambda, u) = e^{a(t) + b(t)\lambda}$. Then,

$$\lambda \partial_t b + \partial_t a + \lambda \int (e^{\delta b z} - 1) e^{u \cdot \bar{z}} d\nu(z) + \lambda \int (e^{u \cdot \bar{z}} - 1) d\nu(z) + \kappa(c - \lambda) b = 0$$

so that

$$\begin{cases} \partial_t b = \kappa b - \int (e^{\delta b z + u \cdot \bar{z}} - 1) d\nu(z) \\ \partial_t a = -\kappa c b \end{cases}$$

with boundary conditions $a(T) = b(T) = 0$. □

Proof of Proposition 4.4. The process $Y = (X, J)^\top$ is Markov since the intensity of J depends only on the current state of X , which is Markov, and the jump distribution of J is fixed. We can append the components of J one at a time. Therefore, it is sufficient to consider the case where J is one dimensional. By assumption, its arrival intensity λ can be expressed as a time-dependent affine function $\lambda(X_t, t) = \Lambda_0(t) + \Lambda_1(t) \cdot X_t$ of X_t .

The first step is to extend the d dimensional jump processes Z^i by one dimension. Suppose the affine point process J is self-affecting so that it is temporally consistent with some Z^i . Renumbering if necessary, assume that $i = 1$ and extend Z^1 by appending the component process J . We retain the symbol ν^1 for the distribution on \mathbb{R}^{d+1} that governs the jump sizes of Z^1 . The extensions of the remaining Z^i 's are made by appending temporally consistent counting processes. In other words, the jumps in the $(d+1)$ st component of Z^2, Z^3, \dots, Z^m are of unit size. It follows that the (degenerate) jump distributions ν^i on \mathbb{R}^{d+1} are canonically determined for $i \geq 2$.

Next, we specify the affine jump diffusion coefficients of Y . Since the intensity λ^i of Z^i is an affine function of X , it can be expressed as an affine function of Y . The one dimensional coefficients Λ_0^i remain the same and the d dimensional vectors Λ_1^i are extended to dimension $(d+1)$ with a zero. The dimension $(d+1)$ sensitivity matrix of Y to Z^1 is given by $\zeta^1 = \text{diag}(0, 0, \dots, 0, 1)$. We extend the $d \times d$ sensitivity matrices $\zeta^2, \zeta^3, \dots, \zeta^m$ by adding one additional row and column with all entries set to zero.

If the affine point process J is not temporally consistent with any Z^i , then all of the X^i 's are extended by counting processes and all the (degenerate) ν^i 's are canonically determined. Here, we add an $(m+1)$ st jump term given by $(d+1)$ copies of J . The intensity $\lambda^{m+1} = \lambda$ so $\Lambda_0^{m+1}(t) = \Lambda_0(t)$ and the $(d+1)$ vector $\Lambda_1^{m+1}(t)$ is gotten from the d vector $\Lambda_1(t)$ by appending a zero. Each of the m sensitivity matrices ζ^i is extended from dimension d to dimension one with a row and column of zeros. The $(d+1)$ dimensional sensitivity matrix $\zeta^{m+1} = \text{diag}(0, 0, \dots, 0, 1)$.

Since J is a pure jump process, the drift coefficients K_0 and K_1 and the covariance matrix coefficients H_0 and H_1 are gotten from the corresponding coefficients for X by extending with rows and columns of zeros. \square

Proposition A.1. *Let J be an affine point process driven by the affine jump diffusion X . Then the conditional transform of $Y = (X, J)^\top$ is given by equation (40) if the coefficient functions β and α uniquely solve the differential equations (41) and (42) and if for $(u, T) \in \mathbb{C}^{d+k} \times \mathbb{R}_+$ the following expectations are all finite:*

- (1) $E(|\Psi_T|)$ where $\Psi_t = \exp(\alpha(t) + \beta(t) \cdot Y_t)$
- (2) $E((\int_0^T \eta_t \cdot \eta_t dt)^{1/2})$ where $\eta_t = \Psi_t \beta(t)^\top \sigma(Y_t, t)$
- (3) $E(\int_0^T |\gamma_t| dt)$ where $\gamma_t = \Psi_t \sum_{i=1}^m (\theta^i(\zeta^i \beta(t)) - 1) \lambda^i(Y_t, t)$.

Proof of Proposition A.1. In view of Proposition 4.4, the result is a corollary of Duffie et al. (2000, Proposition 1). The conditions (1)–(3) guarantee that Ψ is a martingale. \square

Proposition A.2. *Let J be an affine point process driven by the affine jump diffusion X . Then the conditional expectation of $e^{u \cdot Y_T} (v \cdot Y_T)$ is given by formula (44) if the coefficient functions β and α uniquely solve the differential equations (41) and (42), the coefficient functions B and A uniquely solve the differential equations (45) and (46), and if for $(u, v, T) \in \mathbb{C}^{d+k} \times \mathbb{R}^{d+k} \times \mathbb{R}_+$ the following expectations are all finite:*

- (1) $E(|\Phi_T|)$ where $\Phi_t = \Psi_t (A(t) + B(t) \cdot Y_t)$
- (2) $E((\int_0^T \bar{\eta}_t \cdot \bar{\eta}_t dt)^{1/2})$ where $\bar{\eta}_t = \Phi_t (\beta(t)^\top + B(t)^\top) \sigma(Y_t, t)$
- (3) $E(\int_0^T |\bar{\gamma}_t| dt)$ where $\bar{\gamma}_t = \sum_{i=1}^m \lambda^i(Y_t, t) (\Phi_t (\theta^i(\zeta^i \beta(t)) - 1) + \Psi_t \nabla \theta^i(\zeta^i \beta(t)) \cdot \zeta^i B(t))$.

Proof of Proposition A.2. Given Proposition 4.4, the result is a corollary of Duffie et al. (2000, Proposition 3). The conditions (1)–(3) guarantee that Φ is a martingale. \square

Corollary A.3. *In Proposition A.2, suppose that $u = 0$. Then A and B are given by*

$$B(t) = v \exp \left(\int_t^T M_1(s) ds \right), \quad A(t) = \int_t^T M_0(s) \cdot B(s) ds$$

where

$$M_1(t) = K_1(t)^\top + \sum_{i=1}^m \Lambda_1^i(t) \otimes \ell^i \zeta^i, \quad M_0(t) = K_0(t) + \sum_{i=1}^m \Lambda_0^i(t) \ell^i \zeta^i.$$

Proof of Corollary A.3. The assumption $u = 0$ implies that $\alpha = \beta = 0$. Therefore, since $\nabla \theta^i(0)$ is equal to the $(d+k)$ -vector ℓ^i of expected losses of the i th jump term, equations (45) and (46) simplify to

$$\begin{aligned} \partial_t B(t) &= -K_1(t)^\top B(t) - \sum_{i=1}^m \Lambda_1^i(t) \ell^i \cdot \zeta^i B(t) \\ &= -K_1(t)^\top B(t) - \sum_{i=1}^m \Lambda_1^i(t) \otimes \ell^i \zeta^i B(t) \\ &= -M_1(t) B(t) \end{aligned}$$

and

$$\begin{aligned} \partial_t A(t) &= -K_0(t) \cdot B(t) - \sum_{i=1}^m \Lambda_0^i(t) \ell^i \cdot \zeta^i B(t) \\ &= -K_0(t) \cdot B(t) - \sum_{i=1}^m \Lambda_0^i(t) \ell^i \zeta^i \cdot B(t) \\ &= -M_0(t) \cdot B(t) \end{aligned}$$

and the result is immediate. □

References

- Altman, Edward, Brooks Brady, Andrea Resti & Andrea Sironi (2005), ‘The link between default and recovery rates: Theory, empirical evidence and implications’, *Journal of Business* **78**(6), 2203–2227.
- Arnsdorf, Matthias & Igor Halperin (2008), BSLP: Markovian bivariate spread-loss model for portfolio credit derivatives. *Journal of Computational Finance*, forthcoming.
- Azizpour, Shahriar & Kay Giesecke (2008), Premia for correlated default risk. Working Paper, Stanford University.

- Basel Commission on Bank Regulation (2004), International convergence on capital measurement and capital standards. BIS.
- Bassamboo, Achal, Sandeep Juneja & Assaf Zeevi (2008), ‘Portfolio credit risk with extremal dependence: Asymptotic analysis and efficient simulation’, *Operations Research* **56**(3), 593–606.
- Bowsher, Clive (2007), ‘Modelling security market events in continuous time: Intensity based, multivariate point process models’, *Journal of Econometrics* **141**, 876–912.
- Brémaud, Pierre (1981), *Point Processes and Queues – Martingale Dynamics*, Springer-Verlag, New York.
- Brigo, Damiano, Andrea Pallavicini & Roberto Torresetti (2006), Calibration of CDO tranches with the dynamical generalized-Poisson loss model. Working Paper.
- Chen, Zhiyong & Paul Glasserman (2008), ‘Fast pricing of basket default swaps’, *Operations Research* **56**(2), 286–303.
- Cheng, Peng & Olivier Scaillet (2007), ‘Linear-quadratic jump-diffusion modeling’, *Mathematical Finance* **17**, 575–598.
- Collin-Dufresne, Pierre, Robert Goldstein & Jean Helwege (2003), Are jumps in corporate bond yields priced? Modeling contagion via the updating of beliefs. Working Paper, Carnegie Mellon University.
- Cont, Rama & Andreea Minca (2008), Extracting portfolio default rates from CDO spreads. Working Paper, Columbia University.
- Daley, Daryl & David Vere-Jones (2003), *An Introduction to the Theory of Point Processes, Volume I*, Springer-Verlag, New York.
- Das, Sanjiv, Darrell Duffie, Nikunj Kapadia & Leandro Saita (2007), ‘Common failings: How corporate defaults are correlated’, *Journal of Finance* **62**, 93–117.
- Davis, Mark & Violet Lo (2001), Modeling default correlation in bond portfolios, in C.Alexander, ed., ‘Mastering Risk Volume 2: Applications’, Prentice Hall, pp. 141–151.
- Ding, Xiaowei, Kay Giesecke & Pascal Tomecek (2006), Time-changed birth processes and multi-name credit derivatives. *Operations Research*, forthcoming.
- Duffie, Darrell, Andreas Eckner, Guillaume Horel & Leandro Saita (2008), Frailty correlated default. *The Journal of Finance*, forthcoming.
- Duffie, Darrell, Damir Filipovic & Walter Schachermayer (2003), ‘Affine processes and applications in finance’, *Annals of Applied Probability* **13**, 984–1053.

- Duffie, Darrell, Jun Pan & Kenneth Singleton (2000), ‘Transform analysis and asset pricing for affine jump-diffusions’, *Econometrica* **68**, 1343–1376.
- Duffie, Darrell & Nicolae Gârleanu (2001), ‘Risk and valuation of collateralized debt obligations’, *Financial Analysts Journal* **57**(1), 41–59.
- Eckner, Andreas (2007), Computational techniques for basic affine models of portfolio credit risk. Working Paper, Stanford University.
- Giesecke, Kay & Lisa Goldberg (2005), A top down approach to multi-name credit. Working Paper, Stanford University.
- Giesecke, Kay & Stefan Weber (2006), ‘Credit contagion and aggregate losses’, *Journal of Economic Dynamics and Control* **30**(5), 741–767.
- Glasserman, Paul & Jingyi Li (2005), ‘Importance sampling for portfolio credit risk’, *Management Science* **51**(11), 1643–1656.
- Glasserman, Paul & Kyoung-kuk Kim (2008), Saddlepoint approximations for affine jump diffusion models. *Journal of Economic Dynamics and Control*, forthcoming.
- Glasserman, Paul, Wanmo Kang & Perwez Shahabuddin (2007), Fast simulation of multifactor portfolio credit risk. *Operations Research*, forthcoming.
- Hawkes, Alan G. (1971), ‘Spectra of some self-exciting and mutually exciting point processes’, *Biometrika* **58**(1), 83–90.
- Hawkes, Alan G. & David Oakes (1974), ‘A cluster process representation of a self-exciting process’, *Journal of Applied Probability* **11**, 493–503.
- Jarrow, Robert A. & Fan Yu (2001), ‘Counterparty risk and the pricing of defaultable securities’, *Journal of Finance* **56**(5), 555–576.
- Jorion, Philippe & Gaiyan Zhang (2006), Good and bad credit contagion: Evidence from credit default swaps. *Journal of Financial Economics*, Forthcoming.
- Kou, Steven & Xianhua Peng (2009), Default clustering and valuation of collateralized debt obligations. Working Paper, Columbia University.
- Lee, Roger W. (2004), ‘Option pricing by transform methods: Extensions, unification, and error control’, *Journal of Computational Finance* **7**(3), 51–86.
- Longstaff, Francis (2008), The subprime credit crisis and contagion in financial markets. Working Paper, UCLA.
- Longstaff, Francis & Arvind Rajan (2008), ‘An empirical analysis of collateralized debt obligations’, *Journal of Finance* **63**(2), 529–563.

Papageorgiou, Evan & Ronnie Sircar (2007), Multiscale intensity models and name grouping for valuation of multi-name credit derivatives. *Applied Mathematical Finance*, forthcoming.

Protter, Philip (2004), *Stochastic Integration and Differential Equations*, Springer-Verlag, New York.

Revuz, Daniel & Marc Yor (2005), *Continuous Martingales and Brownian Motion*, Springer-Verlag, Heidelberg.