

AA214: NUMERICAL METHODS FOR COMPRESSIBLE FLOWS

Pseudo-Time Integration



Outline

- 1 Vector Form of the Semi-Discrete Euler Equations
- 2 Steady-State Solution
 - Fictitious Time-Evolution
 - Local Time-Stepping



└ Vector Form of the Semi-Discrete Euler Equations

- From Chapter 7, it follows that the semi-discretization on unstructured (or for that matter structured) grids of the unsteady Euler equations in multiple dimensions can be written as

$$\frac{d\bar{W}_i}{dt} = - \sum_{\star} \frac{\hat{\mathcal{F}}_{i\star}}{\|C_i\|}, \quad i = 1, 2, \dots, N$$

where N denotes the total number of finite volume cells

- An alternative expression of the above equations is

$$\|C_i\| \frac{d\bar{W}_i}{dt} + \sum_{\star} \hat{\mathcal{F}}_{i\star} = 0, \quad i = 1, 2, \dots, N$$



Vector Form of the Semi-Discrete Euler Equations

- Let

$$W = [\overline{W}_1^T \ \overline{W}_2^T \ \cdots \ \overline{W}_N^T]^T$$

and

$$F(W) = \left[\sum_{\star} \hat{\mathcal{F}}_{1\star}^T \ \sum_{\star} \hat{\mathcal{F}}_{2\star}^T \ \cdots \ \sum_{\star} \hat{\mathcal{F}}_{N\star}^T \right]^T$$

- And let $\Omega = \text{diag}(\Omega_1, \Omega_2, \dots, \Omega_N)$ denote the diagonal matrix of cell volumes, where $\Omega_i = \|C_i\| \mathbb{I}_m$ ($m = 3, 4, 5$ for one-, two-, and three-dimensional problems, respectively)
- Then, the semi-discretization in multiple dimensions of the unsteady Euler equations can be written in vector form as

$$\Omega \frac{dW}{dt} + F(W) = 0$$



└ Steady-State Solution

└ Fictitious Time-Evolution

- At the steady-state, the above semi-discrete equation simplifies to

$$F(W) = 0$$

- However, because $F(W)$ is in general a highly nonlinear function of W whose solution by Newton's method is *difficult to initialize*, the above nonlinear algebraic equation is transformed into the following nonlinear ordinary differential equation

$$U \frac{dW}{d\tau} + F(W) = 0$$

where U is a *positive definite* matrix, and τ is a *fictitious, global* time

- If the above problem is hyperbolic everywhere, then, regardless of the initial condition, as long as U is positive definite, the above system converges to a steady-state solution as $\tau \rightarrow \infty$
- In particular, the specific choice of a positive definite matrix U is unimportant, except that it may change the convergence point if there are multiple steady-state solutions for W , which is assumed here not to be the case



- └ Steady-State Solution

- └ Local Time-Stepping

$$U \frac{dW}{d\tau} + F(W) = 0$$

- Assume that a different *local* time τ_i is chosen for each cell C_i
- Then, there exists a Jacobian matrix J_i for the transformation from τ in every cell C_i to τ_i : $J_i = \frac{d\tau}{d\tau_i} \mathbb{I}_m$
- Let $J = \text{diag}(J_1, J_2, \dots, J_N)$
- Note that both Ω and J are positive definite matrices
- Assume next that U is chosen as

$$\begin{aligned} U = \Omega J &= \text{diag}(\Omega_1 J_1, \Omega_2 J_2, \dots, \Omega_N J_N) \\ &= \text{diag}(\|C_1\| \frac{d\tau}{d\tau_1} \mathbb{I}_m, \|C_2\| \frac{d\tau}{d\tau_2} \mathbb{I}_m, \dots, \|C_N\| \frac{d\tau}{d\tau_N} \mathbb{I}_m) \end{aligned}$$

- Then, the above equation becomes

$$\Omega J \frac{dW}{d\tau} + F(W) = 0$$



- Steady-State Solution

- Local Time-Stepping

$$\Omega J \frac{dW}{d\tau} + F(W) = 0$$

- The above equation can be re-written as

$$\begin{aligned} \|C_i\| J_i \frac{dW_i}{d\tau} + \sum_{\star} \hat{\mathcal{F}}_{i\star}(W) &= \|C_i\| \frac{d\tau}{d\tau_i} \frac{dW_i}{d\tau} + \sum_{\star} \hat{\mathcal{F}}_{i\star}(W) \\ &= \|C_i\| \frac{dW_i}{d\tau_i} + \sum_{\star} \hat{\mathcal{F}}_{i\star}(W) \\ &= 0, \quad i = 1, 2, \dots, N \end{aligned}$$

- It can also be expressed as

$$\frac{dW_i}{d\bar{\tau}_i} + \sum_{\star} \hat{\mathcal{F}}_{i\star}(W) = 0, \quad i = 1, 2, \dots, N$$

where

$$\bar{\tau}_i = \frac{\tau_i}{\|C_i\|}$$



- Steady-State Solution

- Local Time-Stepping

$$\frac{dW_i}{d\bar{\tau}_i} + \sum_{\star} \hat{\mathcal{F}}_{i\star}(W) = 0, \quad \bar{\tau}_i = \frac{\tau_i}{\|C_i\|}, \quad i = 1, 2, \dots, N$$

- The above equation suggests that the original semi-discrete equation

$$\Omega \frac{dW}{dt} + F(W) = 0$$

can be solved using any preferred ordinary differential equation solver and a *local* time-step



- Steady-State Solution

- Local Time-Stepping

- To understand the benefit of a local time-step, consider the case of a 1D linear advection equation (speed = a) and a FT or BT approximation
- In this case, for a given CFL number CFL , the local and global time-steps are given by

$$\Delta t_i^{\ell} = \frac{CFL}{a} \Delta x_i \quad \text{and} \quad \Delta t_i^g = \frac{CFL}{a} \Delta x_{min}$$

respectively

- It follows that

$$\Delta \bar{t}_i^{\ell} = \frac{\Delta t_i^{\ell}}{\Delta x_i} = \frac{CFL}{a} \quad \text{and} \quad \Delta \bar{t}_i^g = \frac{\Delta t_i^g}{\Delta x_i} = \frac{CFL}{a} \frac{\Delta x_{min}}{\Delta x_i}$$



- Steady-State Solution

- Local Time-Stepping

- Note that

$$\Delta \bar{t}_i^\ell = \frac{\Delta t_i^\ell}{\Delta x_i} = \frac{CFL}{a} \quad \text{and} \quad \Delta \bar{t}_i^g = \frac{\Delta t_i^g}{\Delta x_i} = \frac{CFL}{a} \frac{\Delta x_{min}}{\Delta x_i}$$

$$\Rightarrow \Delta \bar{t}_i^\ell = \Delta \bar{t}_i^g \frac{\Delta x_i}{\Delta x_{min}}$$

- Hence, time-integrating the governing semi-discrete equations using a local time-step Δt_i^ℓ — a process also known as pseudo-time integration — advances the solution in each cell towards the steady-state at the same *scaled* pace
- Comparatively, time-integrating the governing semi-discrete equations using a global time-step Δt_i^g — as in the genuinely unsteady case — slows down convergence toward the steady-state solution
- What happens in the case of a 1D nonlinear advection equation?

