

The Great Recession and Mother's Health

Janet Currie¹

Princeton University

Valentina Duque

Columbia University

Irwin Garfinkel

Columbia University

October 2013

Abstract

We investigate the impacts of the Great Recession on the health of women with children using the last two waves of the Fragile Families and Child Well-being Study. We model the financial crisis with changes in the state unemployment rate and we focus on a wide range of physical and mental health outcomes, as well as health behaviors. Our findings from the individual fixed effects models suggest heterogeneous impacts across demographic and socioeconomic groups. While a rise in the UR worsened the physical and mental health, and increased the likelihood of smoking and using drugs for disadvantaged women (minorities, unmarried, and those with low education), the crisis may have actually improved the health of more advantaged ones (Whites, marrieds, and high skilled). Our results confirm the importance of controlling for individual fixed effects to identify the causal impact of the unemployment on individual health outcomes.

¹ jcurrie@princeton.edu, vd2220@columbia.edu, ig3@columbia.edu.

The authors would like to thank the participants at the Fragile Families Seminar at Columbia University for comments and suggestions. All errors are our own.

I. Introduction

The Great Recession was deeper and longer than any previous recession since the 1930s. From peak - December 2007- to trough -June 2009-, output contracted by 4%, employment rate fell by 6.3%, and the unemployment rate went from 4.8% in April 2008 to 10.6% at its peak in January 2010 (NBER, at <http://www.nber.org/cycles.html>). As of June 2013, the unemployment rate was still 2.8 percentage points above what it was at the start of the recession, the labor force participation rate is 63.5%, the lowest rate since 1978, and the percentage of the population with a job, 58.7%, was stuck near levels last seen in the early 1980s (BLS, 2013; Center for Budget Policy and Priorities, 2013).

The start of the Great Recession was severe, sudden, and sharp, and almost everyone experienced some form of financial, psychological, or physical strain. Recent evidence on the effects of the Great Recession has confirmed that losses were disproportionately concentrated among minorities, youth, low income, and low-education workers (Hoynes, Miller, & Schaller, 2012; Grusky, Western, & Wimer, 2009), and that while men have faced higher unemployment than women, their employment recovery has been faster (Kochhar, 2011).

The Great Recession represented a huge financial and psychological shock for many households, and as such, it may have had a significant impact on health. While a number of studies have examined the relationship between economic downturns and health outcomes, the conclusions are mixed. This study aims to contribute to this discussion by investigating the impacts of the Great Recession on the physical and mental health and health behaviors of women with children. Our study improves on previous research in several ways. First, we are the first to employ a U.S. longitudinal dataset, the Fragile Families and Child Well-being Study (FF), to analyze the impacts of the financial crisis. The longitudinal structure of this dataset allows us to

observe the same woman before and during/after the financial crisis, so we are able to control for individual time-invariant characteristics that might be correlated with both the probability of residing in an area with high unemployment rate and with experiencing declines in health. Second, we focus on a group of disadvantaged women with children, and this is new in the literature. Most evidence regarding the link between economic fluctuations and health has focused on employed workers (usually men), who have the strongest labor force attachment, and hence, little is known about the impacts of the crisis on other groups who have varying degrees of labor force participation but may also be impacted by high unemployment in their communities. Third, by focusing on the period of the Great Recession, we are able to exploit the vast exogenous variation in unemployment rate across states and years, whereas most existing studies using pre-Great Recession data have examined macroeconomic downturns that exhibit less variation and shorter durations. Fourth, the FF provides a wide range of health outcomes – measures of physical and mental health as well as health behaviors – with sufficient variation in each, which allows for an examination of both direct and indirect health responses. Fifth, by exploiting the city level variation in FF, we also show that using a more disaggregated analysis – at the city level --, provides very similar estimates to those obtained from using the state level unemployment rate on women's health outcomes.

Our findings demonstrate the importance of controlling for individual fixed effects when trying to identify the impact of economic fluctuations on health outcomes. Moreover, we claim that the use of very different samples to estimate the effects of economic fluctuations on health, contributes to explaining why previous research has found inconsistent and sometimes null impacts.

Results show sharp differences across demographic and socioeconomic groups. While we find that the crisis worsened the health of more disadvantaged women -- minorities, unmarried, and those with low education --, other groups actually faced better health in some respects. For instance we find that, Blacks and Hispanics, unmarrieds, and mothers with a high school degree or less education were less likely to have “excellent” or “very good” health status or to have good mental health, and that they were more likely to increase health compromising behaviors such as smoking and drug use. Whites, married, and highly educated mothers on the other hand, were likely to have better physical and mental health as unemployment increased. We contrast our findings with estimates obtained from replicating our analysis using the Behavioral and Risk Factor Surveillance System (BRFSS), a nationally representative survey that has been widely used in previous studies. The results show very different patterns. We argue that this could be due to the fact that the BRFSS sample of women is a more advantaged group compared to those in FF – even after constructing a sample of women in BRFSS that was close in observable characteristics to those in FF.

This paper is organized as follows: section II presents a summary of the related literature, section III describes the data, section IV presents the empirical strategy, followed by the results in section V and some extensions in section VI. Section VII provides a brief conclusion.

II. Background

Previous studies of the relationship between economic shocks and their effects on health have come to very different conclusions. A number of studies claim that health improves during economic downturns, arguing that people change their health behaviors (e.g., smoke and drink less, lose weight, practice more exercise, etc.) in response to changes in the economy (i.e., lower wages, changes in time-use, etc.) (e.g., Ruhm, 2000, 2003, 2005; Ruhm & Black, 2002; Dehejia

& Lleras-Muney, 2004). Other studies find that unemployment is associated with poorer health outcomes, a finding that is attributed to the stress associated with losing a job and/or reductions in income and wealth (Dee, 2001; Sullivan & Wachter, 2009; Eliason & Storrie, 2009a, 200b; Browning & Heinesen, 2012). The lack of consensus on how macroeconomic fluctuations are related to changes in health and health behaviors suggests that additional research is warranted.

Table 1 provides an overview of the previous literature about the effects of unemployment on health. As the table shows, most of the previous work has focused on the US and has employed a reduced-form approach that exploits state and year variation in the unemployment rate to examine changes in health and health behaviors. Previous studies have also concentrated on analyzing health outcomes for working-age individuals with the strongest labor force attachment, usually men. Thus, little is known about the impacts of unemployment on other demographic groups (e.g., women, children). All previous U.S. studies have employed cross-sectional data, and most predate the Great Recession. These studies have often used the Behavioral and Risk Factor Surveillance System, which we will use for comparison with our Fragile Families estimates.

a) How Could the Unemployment Rate Affect Health?

In his pioneering study, Ruhm (2000) linked data on state unemployment rates with state-level Vital Statistics Death Records from 1971 to 1992, to examine the link between economic downturns and mortality. He found that, a 1 percentage point increase in the state unemployment rate was associated with a 0.5% reduction in state mortality rates, and he claimed that this result was mainly driven by men of working age.² A large number of subsequent studies have investigated the relationship between economic downturns and health. Ruhm (2003), Neumayer

² In particular, he found significant associations for men between 20 and 44, and weak (null) results for those between 45 and 64 and for those over 65.

(2004), and Gerdtham and Ruhm (2006), using data from the US, Germany, and OECD countries, respectively, confirmed the finding that higher unemployment is associated with lower mortality, and they provided further evidence that individuals are less healthy during economic expansions.³ Dehejia and Lleras-Muney (2004) also found that infant health, measured by the incidence of low birth weight and congenital malformations, improves in times of high unemployment.⁴ These studies attribute the positive link between economic downturns and health to recession-induced changes in health behaviors. They argue that when the economy deteriorates, unemployment rises and wages go down, leading to both income and substitution effects that may affect health through impacts on the demand for medical care, food, and other goods and services such as cigarettes, alcohol, and physical activity (as predicted by the model of “health production” (Grossman, 1972)).

A recent article by Miller et al., (2011) contradicts Ruhm (2000). The authors suggest that the cyclical changes in mortality are concentrated in the young and old, and so are unlikely to represent changes in health behaviors among working age adults. They suggest that the pathway connecting changes in unemployment with changes in mortality has to do with changes in health care quality for individuals 65 and over.

Sullivan and Wachter (2009) followed a large sample of individuals subjected to mass layoffs in Pennsylvania and found significantly higher mortality due to accidents and heart conditions. Eliason and Storrie (2009a), using data from Sweden in 1987 and 1988, examined the effect of plant closings and found a twofold short-term increase in suicides and alcohol-related mortality and a 44% increase in mortality risk among men. Eliason and Storrie (2009b) provided

³ Ruhm (2003) shows that economic recessions are associated with lower incidence of chronic and acute health conditions such as ischemic heart disease, intervertebral disk disorders, and fewer days spent in bed.

⁴ The authors found that Black women are less likely to give birth during recessions (which might be due either to fetal selection or to selection into pregnancy), which tends to raise mean birth weights.

further evidence that job loss significantly increased the risk of hospitalization due to alcohol-related conditions, severe cardiovascular diseases such as myocardial infarction or stroke among men and women, and due to traffic accidents and self-harm among men only. Browning and Heinesen (2012), using Danish administrative data, also found that job loss increases the risk of overall mortality and of death specifically from circulatory disease and traffic accidents.

The disconnect between the literature on the effects of individual job loss and the literature on the effects of unemployment on state-level outcomes suggests that it may be fruitful to investigate the effects of unemployment using individual-level longitudinal data, as we do in this paper.

b) How Could the Unemployment Rate Affect Health-Behaviors?

Studies examining the association between economic fluctuations and health behaviors have generally focused on health-compromising behaviors such as alcohol use and abuse, and cigarette smoking, and have provided mixed evidence. While Ruhm (1995), Ruhm and Black (2002) found that drinking is pro-cyclical, Ruhm (2000) later found that the association between the state unemployment rate and binge drinking is actually positive, although non-significant and Tekin, McClellan, and Minyard (2013) found a negligible impact. Charles and DeCicca (2008), Xu and Kaestner (2010), Dee (2001), and Deb et al., (2011) provided evidence that drinking increases significantly during recessions⁵.

The evidence on smoking is far less abundant, and the results are also uncertain. While a group of studies have found that when the economy contracts, smoking declines (Ruhm, 2000, 2005; Xu & Kaestner, 2010), others claim that smoking is counter-cyclical (Dehejia & Lleras-

⁵ While Charles and DeCicca obtained their result from a sample of young men with high socioeconomic status, Deb and authors confirmed this finding on a sample of workers near retirement age.

Muney, 2004),⁶ or that there is a differential impact across demographic groups. Charles and DeCicca (2008) found that smoking is counter-cyclical for the 10% of men most vulnerable to unemployment (minority and low educated), whereas for those most likely to be employed, smoking falls in times of high unemployment. For the majority of men, unemployment rates were not associated with smoking.

Studies of the effect of economic fluctuations on obesity –a well-established risk factor for cardiovascular disease, high blood pressure, and diabetes – provide similarly ambiguous findings. While Ruhm (2000, 2005) showed that during recessions body mass index (BMI) declines significantly and is particularly driven by those with severe obesity, Charles and DeCicca (2008) and Deb et al., (2011) found an increase in obesity that seems to be driven by minority and low educated groups⁷.

Another important mechanism by which economic fluctuations can affect people's health, regardless of their employment status, is stress. The fear of losing a job, or the actual job loss, can make people anxious or depressed. Studies have found a strong correlation between individual job loss and clinical and subclinical depression, anxiety, and substance use (Murphy & Athanasou, 1999). The evidence in economic studies points out that as the unemployment rate rises, mental health worsens, and this is evident in outcomes such as suicides and suicide attempts (Ruhm, 2000; Browning & Heinesen, 2012); anxiety, depression, loss of confidence and self-esteem (Theodossiou, 1997); feelings of sadness, hopelessness, worthlessness, restlessness,

⁶ Dehejia and Lleras-Muney (2004) found that the positive association between unemployment and smoking is only significant for White mothers (not for Blacks mothers who are positively selected into fertility, particularly in terms of education). When they split the sample by education groups, they found that less educated women with high unemployment see large improvements in health and behavior, whereas health and behavior appear to worsen as education increases.

⁷ Charles and DeCicca find this result on men with low employment probabilities and Deb et al., (2011) suggest that this effect might be driven by the youngest, those with low non-housing net worth, higher depressive symptoms, less education, and females.

and nervousness (Charles & DeCicca, 2008); and substance use (Dee, 2001)⁸. Nevertheless, Tekin, McClellan, and Minyard (2013) find that only better educated individuals experience more mental health problems, and they argue that economic deterioration exacts a larger toll on individuals who have a higher opportunity cost of job loss. Other studies have shown that experiencing unemployment is more strongly associated with mental health problems in men than in women, which could be explained by gender differences in occupation and in family responsibilities -- the nurturing role of women can act as a buffer--. Moreover, differences across marital status show that being married appears to have a protective effect on women. The association between unemployment and mental health problems is larger for single women than for marrieds (Artazcoz et al., 2004).⁹

As this summary indicates, the evidence on the relationship between economic fluctuations and health is far from clear. Countervailing mechanisms and the use of very different samples to estimate these effects make the net impact of economic fluctuations on people's health difficult to predict. In this study we aim to add to this debate by providing further evidence on the effect of the sudden increase in unemployment caused by the Great Recession, on women's physical and mental health, and on their health-behaviors.

Our study provides several contributions relative to the previous literature. First, we use a longitudinal household survey that follows the same mother before and during/after the Great Recession and thus account for time-invariant factors that might be correlated with both the probability of exposure to high unemployment rate and with reporting worse health outcomes.

⁸ Turner (1995) finds some evidence that the effect of individual unemployment on depression and self-reported health status need to be considered within the contextual unemployment rate. The author argues that under low unemployment rate, the association between individual job loss and depression is stronger, particularly among those with a college degree. One caveat to this study is that the author links individual unemployment with individual health measures that are subject to omitted variable bias.

⁹ The evidence shown in this study is not causal; the authors employ multivariate analysis to estimate the association between experiencing individual unemployment and mental health problems, and they analyze differences in these associations by groups. The study uses data for Spain.

Second, by focusing on a group of women with children, we provide new evidence on how the crisis has affected the health of individuals who have varying degrees of labor force participation but might also be significantly impacted by the huge rise in unemployment. Third, we exploit the period of the Great Recession that provides large exogenous variation in labor market conditions across states and years, whereas many of the previous studies have used pre-Great Recession data that exhibit less variation and shorter duration of economic cycles. Fourth, the FF provides rich information of women's health status in several respects –physical and mental health as well as health behaviors –, which allows for an examination of both direct and indirect health responses associated with the crisis. Fifth, we show evidence that using a more disaggregated analysis at the geographic level – using the unemployment rate at the city level --, provides very similar estimates to those obtained from using state level conditions on women's health outcomes.

III. Data

To investigate the effects of the Great Recession on mother's health, we employ the Fragile Families and Child Wellbeing Study (FF), a longitudinal study of 4,897 births that occurred in 20 large U.S. cities located in 15 states, between 1998 and 2000. Unmarried couples were oversampled and constitute about three fourths of the data. When weighted, the sample is representative of urban births in cities with populations over 200,000. Mothers and fathers were interviewed in the hospital shortly after the birth of the focal child, and follow-up interviews were conducted when the focal child was approximately 1, 3, 5, and 9 years old (waves 2, 3, 4, and 5 respectively).

The FF data are uniquely suited to looking at the effects of the Great Recession, as the most recent data collection, year 9, occurred between May 2007 and February 2010. We pooled

years 5 and 9 (periods 2003-2005 and 2007-2010, respectively), which are the years before and during/after the Great Recession. Of the almost 5,000 mothers interviewed at baseline, 87% were interviewed at year 5, and 76% at year 9. We focus on these two waves of data (and exclude years 1 and 3) for two reasons. First, we are interested in examining the impacts of the Great Recession on mother's health. Second, not all health outcomes were available for all waves¹⁰. After restricting the sample to these two waves and to women with complete information on the outcome variables, our analytic sample includes approximately 3,500 mothers (the N varies by the outcome measured).

The outcomes of interest for this study include eight measures of maternal physical and mental health, and health behaviors that were obtained from telephone or in-home interviews, and refer to the last 12 months.¹¹ The following list describes each of the outcomes we investigate. All measures were constructed as binary indicators that take the value of 1 when the mother reports that she has a given condition and 0 otherwise:

Physical health:

- 1) *Self-rated health status*: “excellent” or “very good” health status versus “good”, “fair”, or “poor”.
- 2) *Health problem that limits work*: has a health problem that limits work or study-related activities versus no problem.
- 3) *Obesity*: mother's BMI is equal to or more than 30 versus BMI less than 30.
- 4) *Health insurance*: covered by either a private or Medicaid versus no insurance.

Health Behaviors:

- 5) *Smokes*: smokes cigarettes versus no smoking in the last month.

¹⁰ For example, all the questions related to maternal anxiety were only available at years 1 and 3.

¹¹ These variables include both commonly used health measures and new outcomes that have not been examined in previous studies, for example the use of drugs or an indicator of being diagnosed with clinical depression.

- 6) *Binge Drinking*: drinks 4 or more glasses of alcohol in one occasion versus less than 4 glasses in 1 occasion or no drinking, in the last year.
- 7) *Use drugs*: uses one or more drugs (includes illegal drugs, sedatives, tranquilizers, amphetamines, or other)¹² “on your own”, versus no drug-use. By “on your own” is meant either without a doctor’s prescription, in larger amounts than prescribed, or for a longer period than prescribed.

Mental health:

- 8) Clinical depression (liberal scale): respondent is assessed as clinically depressed¹³ versus not clinically depressed.

Mother Characteristics and Health outcomes

Table 2 presents weighted summary statistics for all the women in the sample, and by race/ethnicity. Descriptive statistics indicate that 62% of the sample report health status that is “excellent” or “very good”, 10% have a problem that limits their work or study activities, a third are obese, and only 81% are covered by a health insurance. In terms of health behaviors, 30% of the mothers smoke, 13% drink more than 4 glasses of alcohol on one occasion (binge drink), and 5% report drug use. Almost 15% of the mothers are assessed as clinically depressed. Compared

¹² The complete list includes: illegal drugs (marihuana or hashish; cocaine or crack or free base; LSD or other hallucinogens; heroin), sedatives (including either barbiturates or sleeping pills such as Seconal, Halcion, Methaqualone), tranquilizers or “nerve pills” (e.g., Librium, Valium, Ativan, Meproamate, Xanax), amphetamines or other stimulants (e.g., methamphetamine, Preludin, Dexedrine, Ritalin, “Speed”), analgesics or other prescription painkillers (note: this does not include normal use of aspirin, Tylenol without codeine, etc., but does include use of Tylenol with codeine and other Rx painkillers like Demerol, Darvon, Percodan, Codeine, Morphine, and Methadone), inhalants (e.g., Amylnitrate, Freon, Nitrous Oxide (“Whippets”), Gasoline, Spray paint).

¹³ A respondent is assessed as clinically depressed (liberal scale) if respondent had feelings of dysphoria (depression) or anhedonia (inability to enjoy what is usually pleasurable) in the past year that lasted for two weeks or more and, if she had symptoms (1.losing interest, 2.feeling tired, 3.change in weight, 4.trouble sleeping, 5.trouble concentrating, 6.feeling worthless, and 7.thinking about death) that lasted at least about half of the day and occurred every day during the two week period. The difference between the liberal and conservative measure of clinically depression is the proportion of the day that respondent feels symptoms (the list of 7 items). The liberal scale considers these symptoms to last "at least about half of the day" (Kessler and Mroczek, 1994, 1997), whereas the conservative scale is "most of the day" (Walters et al., 2002). The conservative scale also includes an additional question (symptom). The liberal scale has been used more often in studies employing the FF panel data.

to national statistics from the Center for Disease Control and Prevention (2012), 55% of women in the US have “excellent” or “very good” health (Whites 65%, Blacks 48%, and Hispanics 51%), 30% are obese, 84.4% are covered by a health insurance coverage, 17% smoke cigarette, 12% binge drink each month, and among mental health problems, 3.7% of women reported feelings of sadness, 2.4% of hopelessness, 2% of worthlessness, and 6% of “everything is an effort” all or most of the time in the past month.¹⁴ These numbers suggest that FF women have on average better or similar physical health to the average US woman, worse mental health, and higher risk health behaviors¹⁵.

Women’s characteristics were all measured at baseline.¹⁶ On average, mothers in FF are 27 at the time of childbirth, a third of the sample is White and a third is Black, 60% have a high school education or less, half are married, more than half are poor or near poor (as shown by an income-to needs ratio that is below 200%), and 50% are employed. The sample is representative of out-of-wedlock births in US urban areas (Reichman et al., 2001).

Differences by race/ethnicity indicate that Whites are more educated, more likely to be married, wealthier, and have a higher probability of being employed than Black or Hispanic mothers. They are also in better physical health (75% have a health status that is excellent or very good and less than a fifth are obese) and better mental health than minority mothers. In terms of substance use, Hispanics and Black mothers are less likely to report that they smoke, drink, and use drugs compared to Whites.

In columns 5 and 6 of Table 2, we split the sample by low versus high unemployment rates with the cutoff being 6% (the average unemployment rate for the period and states of

¹⁴ Statistics obtained from the Center for Disease Control and Prevention for year 2011 (http://www.cdc.gov/nchs/data/series/sr_10/sr10_256.pdf).

¹⁵ No information was available on drug use at the national level.

¹⁶ We use baseline city weights for FF.

interest in this study). The most striking difference between mothers living in areas with high unemployment, and other mothers, is that they are less likely to be White and more likely to be immigrant and poorer. They are also more likely to suffer from obesity and less likely to have health insurance compared to other mothers. These differences in the raw data point to the importance of controlling for differences in the baseline characteristics of mothers in different areas in order to identify the effects of unemployment on health outcomes.

Economic conditions: State Unemployment rate

We obtain data on the state unemployment rate from the Bureau of Labor Statistics' Local Area Unemployment Statistics (LAUS)¹⁷. We construct an average unemployment rate (UR) over the year since the date of a mother's interview, in order to match our key dependent variable which is a health measure over the previous year. The UR is appended to the data based on a mother's baseline state of residence (state in which she was initially sampled at child's birth) and her date of interview, for both years 5 and 9. We use the state in which she was initially sampled to control for the possibility of endogenous migration in response to changes in unemployment rates. Figure 1 shows the large variation in the unemployment rate in all 15 baseline states included in FF for the period 2000 to 2010, and in particular from 2007 when the Great Recession started.¹⁸

In addition to the state unemployment rate, we also exploit the city level geographic variation in FF to conduct a more accurate analysis of the impacts of the Great Recession on women's health. Using the LAUS data, we construct a measure of the average unemployment rate (UR) in the mother's original baseline city (city in which she was initially sampled) and we

¹⁷ The state unemployment and employment-to-population ratio are obtained from Table 3 in the files: "Regional and State Employment and Unemployment" (in pdf format) that are available for each year/month from Dec/1993 to Apr/2012. The link to the BLS with these specific files is: http://www.bls.gov/schedule/archives/laus_nr.htm#2004.

¹⁸ Figure 1 only includes information for the periods of interview.

append it to the FF data based on her Core Based Statistical Area (CBSA, similar to a Metropolitan Statistical Area) and date of the interview.

Control Variables

In models without maternal fixed effects, we include a number of basic socioeconomic and demographic characteristics of the mother that were measured at baseline. These measures include dummy variables for mother's age (<19, 20-23, 24-27, 28-32, 33+), race/ethnicity (White, Black, Hispanic, and other race/ethnicity), education (less than high school, high school, some college, and college or more), immigrant status, marital/relationship status (married, cohabiting, and single), income (we use four categories of income-to-needs ratio¹⁹: poor is less than 100%, near poor income 100-199%, middle income 200-399% and high income 400%+), and child's age (in months). In models with maternal fixed effects all of these variables are already controlled since they are time invariant.

IV. Methods

We estimate the effect of the UR on mother's health using two logistic models, one that pools data from years 5 and 9 and controls for a rich set of covariates and year and state fixed effects, and a second one that accounts for time-invariant mother fixed effects. The following equation describes the first model:

$$Y_{i,t} = \beta_0 + \beta_1 UR_{i,t} + \beta_2 X_{i,t-1} + \alpha_s + \alpha_t + \varepsilon_{i,t} \quad (1)$$

where $Y_{i,t}$ denotes mother i 's health outcome measured at time t , UR the average unemployment rate at baseline state s over the last year t from the date of interview, X is a vector of mother characteristics measured at baseline (described above), and α_s and α_t are vectors of dummies for baseline state and year, respectively. The baseline state dummies control for any time-invariant

¹⁹ The income-to-needs ratio is based on the official U.S. poverty thresholds established by the Census Bureau.

state level factors that are correlated with both state economic conditions and women's health. The year dummies will absorb year specific factors that could affect both the economy and mother's health. ε is the disturbance term. All models are clustered at the baseline state level to account for the within-state correlation in the observations. The coefficient of interest is β_1 .

The second logistic model that controls for mother-specific fixed effects is estimated using equation 2. The only covariate included in this model is α_t , the interview year dummy.

$$Y_{i,t} = \beta_0 + \beta_1 UR_{i,t} + \alpha_t + \varepsilon_{i,t} \quad (2)$$

This model exploits the longitudinal nature of FF to control for observed and unobserved time-invariant characteristics of the mother, that may be correlated with both residing in a state with high UR and experiencing health problems. For instance, if a mother belongs to a demographic group that is likely to be particularly impacted by unemployment, she may also be more likely to suffer from various health problems.

We estimate separate logistic-fixed effects models by subgroups. We stratify the sample by White, Blacks, and Hispanics; marrieds versus unmarrieds; and mothers with a high school degree or less versus those with more than a high school degree. We do these separate analyses because we expect to find heterogeneous impacts across groups, and in particular we hypothesize that the most disadvantaged women (minorities, unmarried, and the least educated) will fare worse during the Great Recession compared to the most advantaged, because they are less likely to be able to insure themselves against contingencies.

V. Results

Table 3 presents results from the pooled Logistic and Logistic-fixed effects models of the impacts of state unemployment rate on health and health behaviors. We only report the

coefficient of interest, β_1 , and when this coefficient is statistically significant at least at the 95 percent level of confidence, we show it in bold type.

The Logistic estimates (equation 1) shown in the first row, with one exception, indicate that health gets worse as unemployment increases, but none of the coefficients are significantly different from zero at the 0.95 level. Self-reported health status and binge drinking are significant at the 0.90 level. Depression, however, is negatively associated with the unemployment rate. While not shown in the regressions due to space limitations, a few covariates are significantly associated with health outcomes in the pooled logistic models. Women with high levels of education report better health outcomes than those with less than a high school degree. Single and cohabiting women have significantly worse health and health behaviors than those who are married. An increase in the income-to-needs ratio measured at baseline is significantly associated with an increase in health status as well as an increase in substance use.

The second row of Table 3 shows models that control for individual fixed effects. These estimates are similar in size to those obtained from the Logistic models, but in a number of cases the coefficients become statistically significant. The estimates confirm that as the economy worsens, women's physical health declines and health compromising behaviors increase. A rise of 1 percentage point in the UR leads to a 15% decline in the odds of having "excellent" or "very good" health, a 26% increase in the odds of smoking, and a 40% rise in the odds of using drugs. Results in Table 3 demonstrate the importance of controlling for individual fixed effects in order to identify the impacts of macroeconomic fluctuations on health outcomes. During the Recession, where the unemployment rate increased from 5 to 10 percent, the likelihood of experiencing poor physical health increased by nearly 75 percent, and the probability of smoking or using drugs increased by more than 100 percent.

No effects are observed on the probability of having health insurance. This result is consistent with Cawley, Moriya, and Simon (2011), who also find no effect on the probability of health insurance for both the working age population and for the sample of women with children.²⁰ Table 3 also shows that the unemployment rate had no effect on obesity.²¹ Previous studies have found mixed evidence on obesity (Ruhm, 2000, 2005; Charles & DeCicca, 2008), however, these studies have not examined the impacts of UR on women. Moreover, no effects were observed on the probability of being diagnosed as clinically depressed.²² Previous research has measured mental health with individual measures of self-rated feelings of sadness, hopelessness, or worthless, or with more extreme measures such as suicides. We use a robust indicator— being diagnosed as clinically depressed – which is based on the World Health Organization Composite International Diagnostic Interview Short Form (CIDI-SF). The CIDI-SF uses information related on a list of different symptoms and their specific duration, and determines a probable diagnosis of the psychiatric condition known as major depressive episode. The CIDI-SF is commonly used in large-scale community surveys (Aalto-Setälä et al., 2002). The fact that we do not observe an effect on clinical depression for the whole sample should not be seen as an inconsistent result with respect to the previous studies. For instance, a recent paper showed that while the 2008 stock market crash – which lead to huge losses in wealth for many households--, caused immediate declines in subjective measures of mental health, it did not increase clinically-validated measures of depressive symptoms or indicators of depression (McInerney, Mellor, & Nicholas, 2013). Moreover, this finding could reflect the fact that the

²⁰ During the recession there was a sharp decline in the share of the population with employer- sponsored health insurance due to the massive decline in employment, which could have been offset by a significant rise in Medicare enrollment (Cawley et al., 2011).

²¹ In another analysis (not shown here) we estimate the effect of UR on the likelihood of being overweight (BMI \geq 25) and we find no effect.

²² In another analysis (not shown here) we also find that a rise in the UR is not associated with changes in the likelihood of being sad or blue for at least two weeks in the past year, which is a less extreme case of mental health condition, providing further evidence that there is little impact of the Recession on the aggregate sample of women.

effects of unemployment on mental health are not equally distributed across groups -- gender, marital status, and education -- (Artazcoz et al., 2004; Tekin et al., 2013).

The bottom panel of Table 3 shows estimates of the UR at the city level²³. Results are highly consistent with those obtained using the state level UR. A 1 percentage point increase in the city unemployment rate is associated with a 12 percent decrease in the likelihood of having “excellent” or “very good” health status (15% decline when we use the state UR), a 30% increase in the odds of smoking (26%), and a 39% rise in the odds of drug use (40%). In the following analysis we present our results using the state UR – rather than the city UR – as it has been more widely used in previous literature and because it provides substantially similar estimates.

Heterogeneous Effects

Results shown in Table 3 assumed that the effects of the Great Recession were born equally by all groups of women. Following the few previous studies that suggest that the effects of the crisis may not have been homogeneously distributed across workers, we examine differences by race/ethnicity, marital status, and education groups. We present only the Logistic-fixed effects models since these provide more reliable estimates.

Table 4 shows the effects of the UR for the whole sample of mothers, and by racial/ethnicity groups (White, Black, and Hispanic), by marital status (married and unmarried), and by education levels (mothers with a high school degree or less education, and those with more than a high school degree). The estimates reveal significant differences in the effects of UR on women’s health across subpopulations, which suggest that more disadvantaged mothers –

²³ Equation 1 is estimated at the city level: it includes the average UR in the last 12 months since the date of interview and in the baseline city, mother covariates measured at baseline, fixed effects at the baseline city and year, and errors are clustered at the baseline city.

minorities, unmarried, and the low educated – were likely to suffer negative health impacts while the most advantaged women experienced some health improvements in certain dimensions.

Results by race/ethnicity indicate that minorities faced larger declines in their physical and mental health compared to Whites. In some respects, the health of Whites improved. For example, a 1 percentage point increase in the UR is predicted to lead to a decline in the likelihood of being obese of almost 50% and a fall in the probability of being diagnosed with clinical depression of 35%. However, it was also associated with an increase in binge drinking of 60%.

The estimates for Blacks (shown in the third row of Table 4) indicate that the odds of having “excellent” or “very good” health fell by 32% and the odds of using drugs rose by 30%. Hispanics also show a marginal increase in drug use and a significant deterioration in mental health (the odds of being diagnosed with clinical depression increase by 30%). This last result contradicts that found in Tekin et al., (2013), in which Hispanic populations show an improvement in mental health when state employment rates decrease.

We now examine differences by marital status. We exploit the fact that the Fragile Families dataset oversamples births to unwed parents, and as a result, it is possible to study differences by mother’s relationship status (married versus unmarried), which is usually difficult to examine in other surveys. These findings indicate that unmarried mothers were hard hit by the Great Recession whereas it may have actually improved the health of marrieds in some respects. Logistic-fixed effects estimates show that a rise in the UR was associated with a decrease in the probability of being depressed among married women (a 40% reduction in the odds), whereas unmarried women were less likely to have “excellent” or “very good” health (19% reduction) and more likely to adopt health compromising behaviors (the odds of smoking rose by a

significant 31% and drug use increased by 29%, though the latter result is not significant at the 95% level).

Finally, results by education group indicate that more educated women suffered less during the Great Recession as they were less likely to have problems that limited their work-related activities (the odds fell by 45%) and were less likely to be clinically depressed (a 23% reduction in the odds, although only significant at the 0.9 level). This result may reflect the fact that those with higher employment prospects (highly educated) suffer less in terms of mental health problems (Charles & DeCicca, 2008). However, more educated women were also more likely to smoke (66%). Less educated mothers on the other hand, faced a significant decline in their physical health as they were more likely to report “good”, “fair”, or “poor” health status (14%) and more likely to have had problems that limited their work (the odds fell by 36%). In terms of health behaviors, the results showed that low educated women faced a significant rise in drug use (41%).

VI. Extensions

Other measures of economic fluctuations: Employment-to-Population Ratio

Unemployed workers who grow discouraged in their job search and do not actively participate in the labor market are not officially counted as unemployed. Hence, reductions in the unemployment rate may sometimes overstate improvements in the labor market. Alternatively, the unemployment rate may remain high even as employment is rising, if discouraged workers come back to the labor market. Thus, in order to more adequately capture fluctuations in the labor market, we also investigate how the state employment-to-population ratio (ER) has affected women’s health. The ER is defined as the number of employed workers as a proportion of the total population aged 18-64 in a given state. We construct an annual average ER measure

using employment data from the LAUS that come from the Current Population Survey, and we append it to FF data based on a mother's baseline state of residence and her date of interview (see foot note 13).

Appendix Table 5 shows Logistic-fixed effects estimates of the effects of state employment-to-population ratio on health outcomes. We find consistent but weaker estimates compared to those obtained when using the state unemployment rate. That is, as the economy expands, women's physical health tends to improve and substance use declines. We also find heterogeneous impacts across demographic groups, indicating that the positive effects of economic recovery (an increase in ER) are mostly experienced by minority, unmarried, and less educated mothers as they are more likely to experience an improvement in their physical and mental health as well as a decrease in the odds of using of drugs. The probability of experiencing clinical depression increases significantly for Whites (43%), marrieds (28%), and highly educated mothers (29%), with better economic conditions, whereas for Hispanics it declines (30%).

Migration

Since almost 20% of the sample has migrated since they were first sampled at childbirth, we also examine how economic conditions in a mother's current state of residence, rather than in her original/baseline state, are associated with her health outcomes. While the measure of UR at the current state captures the labor market conditions in the place of residence for both movers and non-movers, it is, however, endogenous. Mothers may choose to migrate from their baseline state due to the UR shock and move to places with better economic conditions (e.g., lower UR), and if those mothers who move are different in observed and unobserved ways to those that stay, this could bias our estimates of the UR on health. For instance, if migrant mothers are healthier

than those who stay, then the effect of the UR in the state of residence on mother's health may potentially overestimate the true relationship.

To examine how economic conditions in the state of residence affect health outcomes, we construct a measure of state average unemployment rate in a mother's current state of residence, and we append it to FF based on her current state and year of interview. So for instance, if a mother who was originally (at baseline) sampled at Florida, decides to move to New York, from year 5 to year 9, then her UR measure at her current state of residence will include the UR in Florida in year 5 and the UR in New York in year 9. Her UR at baseline will report the UR in Florida in both years 5 and 9.

Appendix Table 6 shows Logistic-fixed effects estimates of the effects of state UR in the current state of residence on mother's health. The estimates show substantially similar coefficients to those obtained when using the baseline UR, which suggests that our estimates of UR on health may not be greatly affected by selective migration. We find that a 1 percentage point increase in the current state unemployment rate is associated with a 15 percent decrease in the likelihood of experiencing "excellent" or "very good" health status (15 percent when we use the baseline/original state UR), and an increase in the probability of smoking and of drug use by 26 percent (22 percent) and 40 percent (36 percent), respectively. Differences by subgroups are also very similar to those shown in Table 4, and confirm our previous finding that more disadvantaged groups of women were more likely to suffer health setbacks during the crisis.

Selective attrition

Another potential source of selection bias in this study is the presence of selective attrition from year 5 to year 9. Selective attrition may bias our estimates of UR on health outcomes if for instance, mothers who are interviewed in year 5 and not in year 9, are missing

from the data perhaps due to experiencing material hardship in year 9 (e.g., telephone service disconnected), which could be correlated with an increase in the probability of experiencing poor health.

The attrition rate from year 5 to year 9 of FF is 19%.²⁴ To analyze whether the effect of UR on the probability of attrition was different for different groups, we perform a simple test in which we construct a dummy variable equal to one if a mother attrited from year 5 to year 9²⁵, and we regress this indicator on the UR she experienced in year 5, her observable characteristics interacted by the UR in year 5, and all other covariates as described in equation 1.²⁶ In the presence of selective attrition, the coefficients on the interaction between the UR and a woman's characteristics should be statistically significant. We also examine selective attrition in terms of health outcomes (physical and mental health and health behaviors) by replicating the previous analysis on selective attrition based on women's observable characteristics, but this time we interact a woman's health outcomes in year 5 with the UR in year 5.

Appendix Tables 11 and 12 show estimates on selective attrition in terms of women's observable characteristics and in terms of women's health outcomes. We find little evidence of selective attrition in terms of mother characteristics. In fact, the only group that is less likely to attrit in year 9 are women with a college degree who faced high UR in year 5. In terms of health status in year 5 we also find little evidence of selective attrition. We do see, however, that women who suffer from obesity in year 5 are significantly more likely to attrit in year 9 after experiencing high unemployment. These findings suggest that selective attrition is not a big issue

²⁴ 3,808 mothers were interviewed in year 5 and 3,069 were interviewed in year 9. These numbers differ slightly from those shown in Table 2 because they are conditional on being interviewed in year 5, hence excludes those mothers who were interviewed in year 9 and not in year 5.

²⁵ The dummy for attrition from year 5 to year 9 takes the value of 1 when the mother was interviewed in year 5 and not in year 9, and zero otherwise.

²⁶ We include a mother's baseline characteristics, and state and year fixed effects. Errors are clustered at the state level.

in our analysis. To the extent that there is selective attrition, women in better health (i.e., more educated, less likely to be obese, and less likely to consume drugs) are more likely to be interviewed in year 9, which may lead to an underestimate of the effect of the UR on women's health.

State UR and Individual labor market outcomes

We now ask how the crisis affected the labor market outcomes of women in our sample, in order to explore the mechanisms underlying the estimated effects on health. Appendix Table 10 shows Logistic-fixed effects estimates of the state UR on women's individual unemployment, employment, and out of the labor force outcomes using equation 2. In general these findings suggest that economic downturns are only weakly associated with the labor market participation of women in FF. Differences by subgroups reveal heterogeneous responses. We find that minorities and unmarried mothers are more likely to be unemployed (or less likely to be employed) as the UR rises. Marrieds in contrast, exhibit a significantly lower probability of unemployment when the economy deteriorates. Although not shown here, we do find stronger associations when we use the state ER, providing more evidence on the weak labor force attachment of this group of women. These results indicate that as the state ER rises, the odds of individual unemployment for all women fall by 10%, and more disadvantaged mothers in particular -- Blacks and Hispanics, unmarried, and low skilled women—are significantly less likely to be unemployed. These results suggest that, other potential mechanisms besides a woman's own exposure to the labor force, may be operating. For instance, having a partner/spouse losing a job could have an effect on household income and/or on maternal stress, which could, in turn, induce changes in health and health behaviors. However, our results suggest that while this channel may be operating for cohabiting women, it does not seem to

affect marrieds, who experience better mental health as the economy deteriorates. One possible explanation is that married women are also more educated, White, and wealthier, and so they are likely to be better cushioned against economic crisis. Two recent articles showed that, unemployment due to layoffs had higher impacts on depression among men than unemployment due to plant closing (Brand, Levy, & Gallo, 2008); and that unemployment decreases the mental health of spouses almost as much as for the directly affected individuals (Marcus, 2013). Hence, since during the Great Recession the probability of unemployment significantly increased for all workers but more for men, minorities and low-skilled, it is likely that the unemployment shock had a lower impact on the mental health of spouses of workers with greater employment prospects (e.g., White, highly educated).

The Effects of Economic Downturns on Health Outcomes in the BRFSS

Previous studies investigating the effects of economic fluctuations on health have often used the Behavioral Risk Factor Surveillance System²⁷ (see Table 1) and have found mixed results. As we discuss here, the BRFSS includes a sample of men and women who are more likely to be advantaged in several socioeconomic dimensions compared to FF women (or to other groups that were likely to be hard hit by the Great Recession). Based on the findings we obtain in this paper – that the negative impacts of the crisis were concentrated only among disadvantaged women --, we hypothesize that women in the BRFSS would be less likely to suffer health setbacks associated with the unemployment shock. Our findings generally support this hypothesis.

The BRFSS is a nationwide telephone survey conducted every year by the Centers for Disease Control and Prevention's Behavioral Surveillance Branch, to measure behavioral risk factors in the adult population (18 years of age or older). The BRFSS is carried out in all 50

²⁷ The BRFSS is publicly available at: http://www.cdc.gov/brfss/annual_data/annual_data.htm.

states and consists of core questions and specific modules included by some states but not others. Most information is comparable over time and across states, and when variables are weighted using BRFSS sampling weights, they produce nationally representative figures. Public health officials use this survey to develop health policies by computing prevalence rates and establishing relationships among variables.

We start by constructing a sample of mothers in BRFSS that is as close as possible to the in FF, for the period of interest – before and during the Great Recession. We select mothers who had a child between 1998 and 2000 and who were living in the center city of a metropolitan statistical area (MSA) in one of the 15 FF baseline states at the moment of interview. Only the most recent years in BRFSS contain some key variables that allow us to identify mothers with these characteristics,²⁸ the final sample includes 4,921 mothers who were living in the center city of an MSA in one of 13 FF baseline states²⁹ for the period 2005-2010.

Appendix Table 9 shows descriptive characteristics for BRFSS and FF mothers. Results indicate that the BRFSS women are a more advantaged group than those in FF, as they are more likely to be White, married, and educated. They are also wealthier.³⁰ Moreover, the cohabiting sample in BRFSS (6%) is very small compared to that in FF (23%). Single mothers in BRFSS are also very different to those in FF since the BRFSS group includes single, divorced, and widowed women. Even among Whites and among Blacks we find striking differences across datasets: these groups are significantly more educated in BRFSS than in FF (this specific result is not shown). In terms of age, mothers in BRFSS are not on average older than those in FF since

²⁸ The key questions that BRFSS asks mothers are: the year of birth of the child and whether the mother lives in the city center of an MSA– these are all important variables for this study as the FF is a national study that follows a cohort of urban children born between 1998 and 2000 --.

²⁹ While FF samples mothers in 15 states, Tennessee and Massachusetts were dropped from the BRFSS sample for not providing sufficient (none) observations on the sample of interest.

³⁰ Even Whites in BRFSS are more educated, more likely to be employed, and less likely to be married/cohabitate than Whites in FF. Also, Blacks in BRFSS are significantly more educated, wealthier, more likely to be employed, and less likely to cohabitate (more likely to be married and single) compared to those in FF.

mother's age in FF is measured at childbirth (1998-2000) and mother's age in BFRSS is measured in the pooled data (2005-2010), hence age is approximately similar in both cases.

We focus on seven measures of self-reported maternal physical and mental health, and health behaviors. We tried to use health measures in BRFSS that could be easily compared to those in FF. One difference across datasets is that the health variables were measured over distinct periods of time. In FF, all health questions refer to the last 12 months whereas in BRFSS they were asked as: "Have you ever...?", making no reference to a specific period³¹. Appendix Table 8 shows the exact questions that were asked to both FF and BRFSS mothers. In general we find that while BRFSS mothers have a self-reported lower health status, they are less likely to be obese, and less likely to smoke and drink. In terms of mental health problems³², BRFSS women are less likely to suffer from them than FF women.

Appendix Table 7 shows estimates of the UR on health outcomes using BRFSS. Next to these estimates we include the coefficients obtained from Logistic-individual fixed effects models using FF (the same coefficients shown in Table 4), to compare findings across datasets. In general, these findings support our hypotheses: the BRFSS results indicate little or no effect of the UR on women's physical and mental health outcomes. Results by demographic groups also show very weak impacts of the UR³³.

Perhaps the most contradictory result between BRFSS and FF is on health behaviors. While in BRFSS women are significantly less likely to smoke (Blacks reduce smoking by 20% and unmarrieds by 14%) and to drink (the odds of binge drinking fall by 6% for all women)

³¹ Only the question related to maternal depression makes reference to the last month: "(Mother) feels stress or depression for 1 day or more in the last month?".

³² It is difficult to compare the mental health conditions of women across datasets since these are measured using different variables (see Appendix Table 8).

³³ We also estimate equivalent models using employment-to-population ratio (instead of the unemployment rate). We find that modeling changes in the economy with the ER did not substantively change our results. These analyses are not reported here but are available from the authors upon request.

when the UR rises, in FF we find that smoking and drinking rise (smoking increases by 26% for all women and Whites and Hispanics increase binge drinking by 60% and 10% respectively). These contrasting results suggest that health behaviors may be quite different in different samples of women. We do find a significant and consistent impact on obesity for Hispanics across surveys: in BRFSS the odds of being obese rise by 9% whereas in FF they increase by 6%.

VII. Discussion and Conclusions

This study contributes to the ongoing discussion of the relationship between economic fluctuations and people's health, by providing new evidence of the effects of the Great Recession on the health of women with children. Our findings show a heterogeneous impact of the UR on women's health. While the recession worsened the health of minorities, unmarried, and low educated mothers, more advantaged women seem to have experienced better health during the Recession, at least in some respects. This result is consistent with recent findings showing that the effects of the crisis were disproportionately concentrated in some subpopulations. In particular, men, Black and Hispanic workers, youth, and those with low education were hardest hit during the Great Recession. Our results imply that the increase in unemployment from 5 to 10% over the two year period 2007-2009, reduced the odds of having "excellent" or "very good" health status by 75% for disadvantaged mothers and increased the odds of smoking and using drugs, or of having serious mental health problems by more than 100%. Wealthier, White, and married mothers had a significant improvement in their physical and mental health, and an increase in smoking and drinking of more than 100%.

While we find changes in health and in health compromising behaviors as the state unemployment increases, the fact that state economic conditions are not a strong predictor of

women's own labor market outcomes (at least when we use the state UR), is indicative that other potential mechanisms, may be driving these effects.³⁴

Previous studies have come to very different conclusions on the impacts of economic downturns on health. Most of this research, however, predates the Great Recession, and only a few of them have focused on the recent crisis. These studies still provide unclear evidence of its consequences on people's health. One potential reason why previous research finds mixed results may be the use of different samples to estimate these effects. We show that while the crisis affected the health of some groups of women, it might have not done so for other groups. We provide further evidence on this hypothesis by showing that the BRFSS women are in general a more advantaged group of women than those in FF, and that the crisis had little effect on their health outcomes

Finally, we show that if we estimate models of the state UR on health outcomes without controlling for individual fixed effects we find small or weak impacts on many outcomes. In contrast, our individual fixed effects specification suggests that economic conditions have significant effects on the health of women, and are particularly harmful to the most disadvantaged women.


³⁴ One possible pathway could operate through the spouse/partner losing a job during the Recession, which in turn could affect household income and/or on maternal stress. However, we do not test this channel of transmission empirically.

VIII. References

- Aalto-Setälä, T., Haarasilta, L., Marttunen, M., Tuulio-Henriksson, A., Poikolainen, K., Aro, H., and Lönnqvist, J. (2002) Major depressive episode among young adults: CIDI-SF versus SCAN consensus diagnosis. *Psychological Medicine*, 32, pp. 1309-1314.
- Artazcoz, L., Benach, J., Borrell, C., and Cortès, I. (2004) Unemployment and Mental Health: Understanding the Interactions Among Gender, Family Roles, and Social Class. *American Journal of Public Health*, Vol 94, No. 1.
- Browning, M. and Heinesen, E. (2012) Effect of Job Loss Due to Plant Closure on Mortality and Hospitalization. *Journal of Health Economics*, 31, 599-616.
- Bureau of Labor Statistics (July, 2013) News Release: The Employment Situation June, 2013. <http://www.bls.gov/news.release/pdf/empsit.pdf>
- Cawley, J., Moriya, A.S. and Simon, K.I. (2011) The Impact of the Macroeconomy on Health Insurance Coverage: Evidence from the Great Recession. *NBER*, # w17600.
- Charles, K.K. and DeCicca, P. (2008) Local labor Market Fluctuations and Health: Is There a Connection and for Whom? *Journal of Health Economics*, 27(6): 1532-1550.
- Center for Budget Policy and Priorities (July, 2013) Chart Book: The Legacy of the Great Recession. Special Series: Economic Recovery Watch. <http://www.cbpp.org/cms/index.cfm?fa=view&id=3252>
- Center for Disease Control and Prevention (December, 2012) Summary Health Statistics for U.S. Adults: National Health Interview Survey, 2011- *Vital and Health Statistics*. Series 10, Number 256. http://www.cdc.gov/nchs/data/series/sr_10/sr10_256.pdf
- Dee, T.S. (2001) Alcohol Abuse and Economic Conditions: Evidence from Repeated Cross-Sections of Individual-Level Data. *Health Economics*, 10: 257-270.
- Deb, P., Gallo, W. T., Ayyagari, P., Fletcher, J., and Sindelar, J. (2011) The Effect of Job Loss on Overweight and Drinking. *Journal of Health Economics*, 30(2):317-327.
- Dehejia, R. and Lleras-Muney, A (2004) Booms, Busts, and Babies' Health. *Quarterly Journal of Economics*, 119(3): 1091-1130.
- Eliason, M. and Storrie, D. (2009a) Does Job Loss Shorten Life? *Journal of Human Resources*, 44(2), 277-302.
- Eliason, M. and Storrie, D. (2009b) Job Loss is Bad for Your Health: Swedish Evidence on Cause-Specific Hospitalization Following Involuntary Job Loss, *Social Science and Medicine*, 68(8), 1396-1406.
- Gerdtham, U.G. and Ruhm, C. (2006) Deaths Rise in Good Economic Times: Evidence from the OECD. *Economics & Human Biology*, 4(3), 298-316.
- Goodman, C.J., and Mance, S.M. (2011) Employment Loss and the 2007–09 Recession: An Overview. *Monthly Labor Review*, 134(4): 3–12.

- Grossman, M. (1972) *The Demand for Health – A theoretical and Empirical Investigation*. New York: *National Bureau of Economic Research*.
- Hoynes, H., Miller, D., and Schaller, J. (2012) Who Suffers During Recessions? *Journal of Economic Perspectives*, 26(3): 27-48.
- Huff-Stevens, A. Miller, D.L., Page, M.E., and Filipowski, M. (2011) The Best of Times, the Worst of Times: Understanding Pro-cyclical Mortality. *NBER*, # w17657.
- Kochhar, R. (2011). Two Years of Economic Recovery: Women Lose Jobs, Men Find Them. *Pew Research Social and Demographic Trends Project*. Washington DC.
- Marcus, J. (2013). The effect of unemployment on the mental health of spouses – Evidence from plant closures in Germany. *Journal of Health Economics*, 32, 546-558.
- McInerney, M., Mellor, J.M., & Nicholas, L.H. (2013) Recession Depression: Mental Health Effects of the 2008 Stock Market Crash. *Journal of Health Economics*, 32, 1090-1104.
- Murphy, G.C. and Athanasou, J.A. (1999) Unemployment and general health: A review and evaluation, *Journal of Occupational and Organisational Psychology*, 72, 83-99.
- Neumayer, E. (2004) Recessions Lower (some) Mortality Rates: Evidence from Germany. *Social Science & Medicine*, 58(6), 1037-1047.
- Reichman, N. Teitler, J., Garfinkel, I., and McLanahan, S. (2001) Fragile Families: Sample and Design. *Children and Youth Services Review*. 23(4/5): 303-326.
- Ruhm, C. (2000) Are Recessions Good for Your Health? *Quarterly Journal of Economics*, 115(2): 617-650.
- Ruhm, C. (2003) Good Times Make You Sick. *Journal of Health Economics*, 22(4): 637-658.
- Ruhm, C. (2005) Healthy living in hard times. *Journal of Health Economics*, 24, 341-363.
- Ruhm, C. and Black, W.E. (2002) Does Drinking Really Decrease in Bad Times? *Journal of Health Economics*, 21(4), 659-678.
- Sullivan, D. and von Wachter, T. (2009) Job Displacement and Mortality: An Analysis using Administrative Data. *Quarterly Journal of Economics*, 1265-1306.
- Tekin, E., McClellan, C., and Minyard, K.J. (2013) Health and Health behaviors During the Worst of Times: Evidence from the Great Recession. *NBER* # w19234.
- Theodossiou, I. (1997) The Effects of Low-Pay and Unemployment on Psychological Well-Being: A Logistic Regression Approach. *Journal of Health Economics*, 17, 85-104.
- Turner, J.B. (1995) Economic Context and the Health Effects of Unemployment. *Journal of Health and Social Behavior*, 36, 213-229.
- Xu, X. and Kaestner, R. (2010) The Business Cycle and Health Behaviors. *NBER*, # w15737.

Figure 1: State Unemployment Rate (%) During Interview



Note: sample includes the 15 baseline states in FF.

Table 1 – Effects of UR on Health Outcomes

<i>OUTCOME VARIABLE</i>						
<u>Study and Data</u>	Health status/ Health problem that limits work	Weight	Use of medical care / Health insurance	Smokes	Drinks	Mental health

1. Studies Examining Men and Women

<p>Ruhm (2000)</p> <p>Micro Data: - BRFSS: 1987–1995 N =114,000 men and women</p> <p>Methods: Linear model</p>		<p>A 1 pp rise in the UR reduces:</p> <ul style="list-style-type: none"> - Underweight by 0.06pp - Overweight by 0.17pp - Obesity by 0.21pp <p>No analysis by gender</p>	<p>A 1 pp rise in the UR reduces:</p> <ul style="list-style-type: none"> - Visits to the doctor by 0.57pp <p>No analysis by gender</p>	<p>A 1 pp rise in the UR:</p> <ul style="list-style-type: none"> - Reduces smoking by 0.3pp - Reduces # of cigs per day by 0.8pp for current smokers <p>No analysis by gender</p>	<p>A 1 pp rise in the UR:</p> <ul style="list-style-type: none"> - Increases alcohol consumption but is NON-significant <p>No analysis by gender</p>	
<p>Ruhm and Black (2002)</p> <p>Micro Data: - BRFSS: 1987-1999 - N = 500,000 men and women</p> <p>Methods: Linear model</p>					<p>A 1 pp rise in UR:</p> <ul style="list-style-type: none"> - Reduces predicted drinking by 0.2pp - Reduces log(# of drinks) in last-month by 3.1pp for drinkers - No effect on binge drinking <p>By gender:</p> <ul style="list-style-type: none"> - No difference in the probability of drinking - Higher decline for men: # of drinks falls by 3.4pp versus 2.8pp for females 	

Study and Data	Health status/ Health problem that limits work	Weight	Use of medical care / Health insurance	Smokes	Drinks	Mental health
<p>Ruhm (2003)</p> <p>Micro Data: - NHIS: 1972–1981 for individuals 30+ 3 samples: 1) Full sample: N =217,471 men and women 2) 30–64-year-old workers: N = 115,463 obs, men and women 3) 30–55-year-old working males: N = 57,633</p> <p>Methods: Linear model</p>	<p>A 1pp fall in the UR: -For everyone: increases the probability of having a medical condition (chronic or acute) by 0.6 pp -Stronger effects for men of working age</p> <p>By gender: - For men 30-55: increases by 0.63pp -For women 30-55: increases by 0.59pp</p>		<p>Hospital visit is NON-significant in all cases</p> <p>A 1pp fall in the UR: -For everyone: the probability of hospitalization falls by 0.11pp and doctor visits rises by 0.3 although NON-significant -Stronger effects for men of working age</p> <p>By gender results were non-significant: - For men 30-55: falls by 0.11pp -For women 30-55: falls by 0.13pp</p>			
<p>Ruhm (2005)</p> <p>Micro Data: - BRFSS: sample includes individuals aged 18+ from the 1987-2000 waves N = approx. 1.5 million obs, men and women</p> <p>Methods: Linear model</p>		<p>-A 1 pp fall in the employment rate: - Reduces obesity by 0.07pp - Reduces severely obese by 0.4pp - No effect on overweight</p> <p>No differences by gender</p>		<p>-A 1 pp fall in the employment rate reduces: - Smoking by 0.13pp - Smoking of at least 20 cigs/day by 0.10pp smoking at least 40 cig/day by 0.015pp</p> <p>By gender: The effect on smoking is stronger for females (0.17pp) than for males (0.09pp)</p>		

Study and Data	Health status/ Health problem that limits work	Weight	Use of medical care / Health insurance	Smokes	Drinks	Mental health
<p>Dee (2001)</p> <p>Micro Data: - BRFSS, 1984-1995 - N > 700,000, 'prime-age' men and women 18+ (average age is 45)</p> <p>Methods: Linear model</p>					<p>A 5pp increase in UR: - Reduces drinks per month by 3.5% and chronic drinking participation (60 or more drinks/month) by 19% - Binge drinking rises by 8%</p> <p>By gender: - The probability of binge drinking for males is >3 times larger than that for females</p>	
<p>Tekin, McClellan, and Minyard (2013)</p> <p>Micro Data: - BRFSS Sample: N=849,594, individuals in the Labor force, ages 25-55</p> <p>Methods: Linear model</p>	<p>A 1pp drop in ER: - Probability of being in poor health rises by 0.00067 pp (4.8%)</p> <p>By gender: - Strongest effects on females: "excellent health" falls by 0.0044pp (1.8%) and poor health increases by 0.0015pp(10%)</p>				<p>A 1pp drop in ER: - Binge drinking (respondent reports had 60 or more drinks during the past month) falls by 0.0023pp</p> <p>By gender: - Effect on binge drinking is driven by women (0.0022pp)</p>	

2. Studies Examining Men Only

Study and Data	Health status/ Health problem that limits work	Weight	Use of medical care / Health insurance	Smokes	Drinks	Mental health
<p>Charles and DeCicca (2008)</p> <p>Micro Data: - NHIS: Sample of working-aged men in the US living in "large" MSAs: 1997-2001 N=30,000-35,000 men of working age</p> <p>Methods: Linear model</p>		<p>A 1 pp rise in the UR: - Men with low ex-ante employment probabilities experience an increase in underweight by 0.5pp, in overweight by 3.5pp, and in obesity by 2.pp</p> <p>NO effects on men with high employment probability</p>		<p>A 1 pp rise in the UR: - Smoking rises by 2.7pp for those in the lowest employment decile - 2.3pp reduction for those most likely to be employed</p>	<p>A 1 pp rise in the UR: - Only for the group with highest employment probability drinking rises (3.6pp) - # of days with 5 drinks or more declines for all BUT it is NON significant</p>	<p>A 1 pp rise in the UR: - Leads to rises in: sadness (1.5pp), hopelessness (1.1pp), worthlessness (0.9pp), restlessness (1.2pp), nervousness (1.3pp), and feelings of effort (2.5pp)</p> <p>- Strongest effects for those least likely to be employed</p>
<p>Xu and Kaestner (2010)</p> <p>Micro Data: - BRFSS (1984- 2005, N=506,753) and NHIS (1976-2001, N=216,113), men between 25-55</p> <p>Methods: Linear model</p>			<p>A 2.5 % increase in employment: - Decreases doctor visits by 1.5pp</p>	<p>A 2.5 % increase in employment: - Increases smoking by 1 pp - Increases smoking intensity (smokes >19 cigarettes a day y/n) by 2pp</p>	<p>A 2.5 % increase in employment: - Increases drinking by 0.1pp - Decreases binge drinking by 0.2pp</p>	

Table 2: Summary Statistics in FF

Variable	Full	Whites	Blacks	Hispanics	UR>6%	UR<=6%
<u>Health outcomes:</u>						
Health Excellent/V. Good	0.612	0.750	0.547	0.530	0.607	0.613
Health Limits Work	0.103	0.072	0.130	0.113	0.106	0.101
Obesity	0.320	0.195	0.402	0.386	0.348	0.302
Health Insurance	0.809	0.910	0.861	0.640	0.793	0.820
Smokes	0.302	0.369	0.338	0.199	0.295	0.307
>=4 Drinks 1 Time Last Yr.	0.126	0.206	0.083	0.108	0.126	0.127
Drugs use	0.053	0.063	0.063	0.039	0.062	0.046
Clinically depressed (liberal scale)	0.132	0.128	0.147	0.130	0.132	0.132
<u>Race/ethnicity:</u>						
White	0.284	1.000			0.265	0.298
Black	0.357		1.000		0.351	0.361
Hispanic	0.290			1.000	0.308	0.276
Other race	0.070				0.077	0.065
Immigrant	0.239	0.073	0.075	0.460	0.263	0.221
<u>Education:</u>						
<HS	0.271	0.116	0.286	0.455	0.284	0.262
HS	0.329	0.219	0.436	0.320	0.327	0.331
Some college	0.203	0.207	0.236	0.173	0.210	0.197
College or >	0.197	0.457	0.042	0.053	0.180	0.211
<u>Marital Status:</u>						
Married	0.519	0.818	0.231	0.502	0.517	0.520
Cohabiting	0.233	0.114	0.297	0.309	0.234	0.233
Single	0.248	0.068	0.472	0.189	0.249	0.247
Age	26.9 (6.2)	30.0 (6.1)	24.9 (5.8)	25.6 (5.7)	25.1 (5.9)	25.1 (6.0)
<u>Income-to-needs ratio:</u>						
<100%	0.254	0.056	0.382	0.338	0.263	0.247
100-199%	0.276	0.161	0.315	0.329	0.297	0.260
200-299%	0.136	0.122	0.152	0.148	0.133	0.138
>=300%	0.334	0.661	0.151	0.185	0.306	0.354
<u>Employment status:</u>						
Employed	0.478	0.573	0.471	0.400	0.469	0.486
Unemployed	0.175	0.070	0.274	0.185	0.171	0.177
Out of Labor Force	0.344	0.357	0.252	0.415	0.356	0.336
<i>N pooled sample</i>	7,080	1,446	3,515	1,873	2,748	4,332
<i>N year 5</i>	3,829	773	1,887	1,035	931	2,898
<i>N year 9</i>	3,251	673	1,628	838	1,817	1,434

Note:

All characteristics are measured at baseline.
Sample includes mothers interviewed in years 5 and 9.
Numbers are weighted using baseline city weights.

Table 3: The Effect of the UR on Mother's Health in FF

	Health status Excellent or Very Good	Health problem limits work	Obesity	Any health insurance	Smokes	>=4 Drinks on 1 Occasion	Drug use	Clinically depressed
State UR								
<i>LOGIT</i>	0.933 [-1.755]	0.989 [-0.284]	1.024 [1.407]	0.987 [-0.334]	1.090 [1.181]	1.091 [1.852]	1.029 [0.471]	0.998 [-0.033]
<i>LOGIT-FE</i>	0.848 [-2.602]	1.073 [0.661]	1.087 [0.712]	0.993 [-0.111]	1.256 [2.231]	1.140 [1.521]	1.400 [2.499]	0.941 [-0.791]
City UR								
<i>LOGIT</i>	0.916 [-1.951]	1.058 [1.365]	1.025 [1.237]	0.956 [-0.971]	1.088 [1.790]	1.062 [1.339]	1.128 [1.576]	0.994 [-0.107]
<i>LOGIT-FE</i>	0.880 [-2.112]	1.154 [1.451]	1.038 [0.332]	1.002 [0.030]	1.295 [2.650]	1.151 [1.640]	1.394 [2.738]	0.938 [-0.851]
N	7,080	7,070	6,178	7,064	7,079	7,070	7,058	7,067
N changers	953	364	511	758	359	491	322	591
Indiv. controls	Y	Y	Y	Y	Y	Y	Y	Y
Bline State / City FE	Y	Y	Y	Y	Y	Y	Y	Y
Year FE	Y	Y	Y	Y	Y	Y	Y	Y

Note:

Each coefficient comes from a separate regression.

Logit models control for mother characteristics (age, race, education, marital status), and state (city) and year fixed effects. Errors are clustered at the baseline state (city) level.

T-statistics are shown in brackets; bold font indicates that the result is statistically significant at the 95% level of confidence.

Table 4: Logit-FE Estimates of Effects of State UR Using FF

	Health status Excellent or Very Good	Health problem limits work	Obesity	Any health insurance	Smokes	>=4 Drinks on 1 Occasion	Drug use	Clinically depressed
All	0.848 [-2.602]	1.073 [0.661]	1.087 [0.712]	0.993 [-0.111]	1.256 [2.231]	1.140 [1.521]	1.400 [2.499]	0.941 [-0.791]
White	0.909 [-0.691]	0.939 [-0.252]	0.508 [-2.091]	1.034 [0.221]	1.512 [1.571]	1.594 [2.535]	1.230 [0.759]	0.663 [-2.402]
Black	0.687 [-3.009]	1.310 [1.532]	1.358 [1.312]	0.848 [-1.165]	1.227 [1.121]	1.148 [0.672]	1.703 [2.456]	0.865 [-1.002]
Hispanic	0.933 [-0.781]	1.157 [0.792]	1.151 [0.850]	1.061 [0.651]	1.060 [0.341]	1.073 [0.564]	1.902 [1.602]	1.301 [1.982]
Married	1.054 [0.342]	0.663 [-1.245]	0.761 [-0.993]	1.064 [0.334]	0.903 [-0.321]	1.079 [0.331]	1.656 [1.340]	0.596 [-2.371]
Unmarried	0.811 [-2.940]	1.146 [1.191]	1.197 [1.331]	0.964 [-0.520]	1.310 [2.462]	1.128 [1.291]	1.292 [1.732]	1.018 [0.211]
More than HS	0.858 [-1.301]	0.566 [-2.274]	1.16 [0.451]	0.872 [-1.147]	1.655 [2.119]	1.116 [0.742]	1.400 [1.291]	0.771 [-1.950]
HS or less	0.856 [-1.971]	1.364 [2.353]	1.127 [0.916]	1.047 [0.589]	1.144 [1.164]	1.162 [1.391]	1.413 [2.144]	1.062 [0.601]
N	7,080	7,070	6,178	7,064	7,079	7,070	7,058	7,067
Indiv. controls	Y	Y	Y	Y	Y	Y	Y	Y
Bline State FE	Y	Y	Y	Y	Y	Y	Y	Y
Year FE	Y	Y	Y	Y	Y	Y	Y	Y

Note:

Each coefficient comes from a separate regression.

Logit models control for mother characteristics (age, race, education, marital status), and state and year fixed effects. Errors are clustered at the baseline state level.

T-statistics are shown in brackets; bold font indicates that the result is statistically significant at the 95% level of confidence.

Appendix

Appendix Table 5: Logit-FE Estimates of Effects of State ER Using FF

	Health status Excellent or Very Good	Health problem limits work	Obesity	Any health insurance	Smokes	>=4 Drinks on 1 Occasion	Drug use	Clinically depressed
All	1.043 [1.164]	0.995 [-0.09]	0.924 [-1.191]	1.023 [0.530]	0.956 [-0.774]	0.919 [-1.365]	0.902 [-1.576]	1.015 [0.341]
White	0.812 [-1.700]	1.342 [1.708]	0.927 [-0.331]	1.030 [0.238]	0.961 [-0.265]	0.800 [-1.862]	0.920 [-0.531]	1.430 [2.611]
Black	1.006 [0.142]	0.967 [-0.492]	0.975 [-0.312]	1.081 [1.396]	0.983 [-0.228]	1.022 [0.261]	0.873 [-1.676]	1.021 [0.352]
Hispanic	1.043 [1.155]	0.995 [-0.094]	0.821 [-1.142]	0.835 [-1.844]	1.044 [0.249]	0.878 [-1.075]	1.009 [0.031]	0.706 [-2.764]
Married	0.932 [-0.857]	1.045 [0.337]	0.783 [-1.292]	0.877 [-1.183]	1.075 [-0.421]	0.843 [-1.261]	-	1.278 [2.004]
Unmarried	1.070 [1.643]	0.990 [-0.165]	0.940 [-0.863]	1.062 [1.281]	0.930 [-1.132]	0.961 [-0.660]	0.874 [-1.772]	0.981 [-0.381]
More than HS	0.963 [-0.588]	1.284 [2.279]	1.006 [0.043]	0.999 [-0.013]	0.995 [-0.04]	0.891 [-1.190]	0.966 [0.281]	1.291 [2.742]
HS or less	1.080 [1.714]	0.899 [-1.599]	0.895 [-1.439]	1.031 [0.621]	0.941 [-0.904]	0.953 [-0.718]	0.871 [0.070]	0.934 [-1.241]
N	7,080	7,070	6,178	7,064	7,079	7,070	7,058	7,067
Indiv. controls	Y	Y	Y	Y	Y	Y	Y	Y
Blind State FE	Y	Y	Y	Y	Y	Y	Y	Y
Year FE	Y	Y	Y	Y	Y	Y	Y	Y

Note:

Each coefficient comes from a separate regression.

Logit models control for mother characteristics (age, race, education, marital status), and state and year fixed effects. Errors are clustered at the baseline state level.

T-statistics are shown in brackets; bold font indicates that the result is statistically significant at the 95% level of confidence.

Appendix Table 6: Logit-FE Estimates of Effects of Current State of Residence UR in Using FF

	Health status Excellent or Very Good	Health problem limits work	Obesity	Any health insurance	Smokes	>=4 Drinks on 1 Occasion	Drug use	Clinically depressed
All	0.853 [-2.644]	1.091 [0.827]	1.073 [0.632]	0.992 [-0.125]	1.220 [2.046]	1.108 [1.251]	1.355 [2.509]	1.019 [0.258]
White	0.931 [-0.442]	1.086 [0.328]	0.824 [-0.722]	1.008 [0.052]	1.459 [1.616]	1.216 [1.152]	1.232 [0.899]	0.767 [-1.593]
Black	0.706 [-3.138]	1.321 [1.644]	1.175 [0.772]	0.933 [-0.558]	1.149 [0.832]	1.276 [1.338]	1.480 [2.134]	0.955 [-0.372]
Hispanic	0.925 [-0.915]	1.067 [0.359]	1.072 [0.434]	1.046 [0.495]	1.076 [0.423]	1.058 [0.455]	1.942 [1.669]	1.339 [2.170]
Married	1.080 [0.547]	0.588 [-1.622]	0.782 [-1.030]	0.983 [-0.097]	1.019 [0.060]	0.951 [-0.220]	1.734 [1.630]	0.715 [-1.700]
Unmarried	0.807 [-3.146]	1.205 [1.609]	1.186 [1.299]	0.968 [-0.477]	1.278 [2.327]	1.120 [1.279]	1.225 [1.508]	1.095 [1.106]
More than HS	0.882 [-1.151]	0.599 [-2.091]	1.347 [1.009]	0.924 [-0.738]	1.359 [1.461]	1.104 [0.677]	1.441 [1.563]	0.881 [-1.016]
HS or less	0.850 [-2.222]	1.319 [2.160]	1.088 [0.673]	1.026 [0.334]	1.155 [1.297]	1.120 [1.100]	1.341 [1.991]	1.122 [1.195]
N	7,080	7,070	6,178	7,064	7,079	7,070	7,058	7,067
Indiv. controls	Y	Y	Y	Y	Y	Y	Y	Y
State FE	Y	Y	Y	Y	Y	Y	Y	Y
Year FE	Y	Y	Y	Y	Y	Y	Y	Y

Note:

Each coefficient comes from a separate regression.

Logit models control for mother characteristics (age, race, education, marital status), and state and year fixed effects.

Errors are clustered at the state level.

T-statistics are shown in brackets; bold font indicates that the result is statistically significant at the 95% level of confidence.

Appendix Table 7: Logit Estimates of Effects of State UR Using BRFSS

	Health status Excellent or Very Good		Health problem limits work		Obesity		Any health insurance		Smokes		>=4 Drinks on 1 Occasion		Depressed ¹	
	BRFSS	FF	BRFSS	FF	BRFSS	FF	BRFSS	FF	BRFSS	FF	BRFSS	FF	BRFSS	FF
All	1.014 [0.507]	0.848 [-2.602]	0.950 [-1.241]	1.073 [0.661]	1.019 [0.450]	1.087 [0.712]	0.950 [-1.191]	0.993 [-0.111]	0.949 [-1.084]	1.256 [2.231]	0.939 [-1.732]	1.140 [1.521]	0.992 [-0.201]	0.941 [-0.791]
White	1.058 [1.482]	0.909 [-0.691]	1.017 [0.312]	0.939 [-0.252]	1.007 [0.226]	0.508 [-2.091]	0.790 [-2.037]	1.034 [0.371]	0.995 [-0.112]	1.512 [1.571]	0.909 [-1.301]	1.594 [2.535]	1.059 [1.131]	0.663 [-2.402]
Black	1.009 [0.082]	0.687 [-3.009]	0.817 [-1.501]	1.310 [1.532]	1.014 [0.203]	1.358 [1.312]	1.071 [0.643]	0.848 [-1.165]	0.807 [-2.523]	1.227 [1.121]	0.938 [-0.992]	1.148 [0.672]	0.872 [-3.152]	0.865 [-1.002]
Hispanic	1.045 [1.033]	0.924 [-1.812]	1.074 [0.923]	1.058 [0.960]	1.090 [4.362]	1.063 [2.567]	1.007 [0.111]	1.001 [0.042]	0.979 [-0.192]	1.066 [1.721]	0.998 [-0.023]	1.102 [2.352]	0.987 [-0.182]	1.170 [4.971]
Married	1.033 [0.084]	1.054 [0.342]	0.996 [-0.003]	0.663 [-1.245]	1.001 [0.037]	0.761 [-0.993]	0.975 [-0.376]	1.064 [0.334]	1.109 [0.990]	0.903 [-0.321]	0.888 [-1.841]	1.079 [0.331]	1.022 [0.571]	0.596 [-2.371]
Unmarried	1.000 [0.00]	0.811 [-2.940]	0.902 [-0.950]	1.146 [1.191]	1.033 [0.484]	1.197 [1.331]	0.921 [-1.281]	0.964 [-0.520]	0.858 [-3.626]	1.310 [2.462]	1.011 [0.158]	1.128 [1.291]	0.952 [-0.762]	1.018 [0.211]
More than HS	1.056 [1.216]	0.858 [-1.301]	0.932 [-0.714]	0.566 [-2.274]	0.961 [-0.652]	1.160 [0.451]	1.032 [0.509]	0.872 [-1.147]	0.944 [-0.741]	1.655 [2.119]	0.912 [-2.001]	1.116 [0.742]	1.1015 [0.382]	0.771 [-1.950]
HS or less	0.982 [-0.442]	0.856 [-1.971]	0.954 [-0.379]	1.364 [2.353]	1.103 [1.759]	1.127 [0.916]	0.911 [-1.273]	1.047 [0.589]	0.966 [-0.551]	1.144 [1.164]	0.99 [-0.102]	1.162 [1.391]	0.948 [-1.012]	1.062 [0.601]
N	4,908	7,080	4,910	7,070	4,583	6,178	4,913	7,064	4,911	7,079	4,321	7,070	4,921	7,067
Indiv. controls	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
State FE	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y
Year FE	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y

Note:

BRFSS results were estimated using a linear model while those from FF were estimated using individual-fixed effects.

Each coefficient comes from a separate regression.

Logit models control for mother characteristics (age, race, education, marital status), and state and year fixed effects. Errors are clustered at the state level.

T-statistics are shown in brackets; bold font indicates that the result is statistically significant at the 95% level of confidence.

¹ In BRFSS the measure for depression is "Depressed, sad, blue for 2-weeks in the last-year", whereas in FF the measure is Clinically depressed.

Appendix Table 8 - Survey Questions in FF and BRFSS

	FF	BRFSS
<u>Physical health</u>		
Health status	In general, how is your health? Would you say it is: A. Excellent B. Very good C. Good D. Fair E. Poor	Would you say that in general your health is: A. Excellent B. Very good C. Good D. Fair E. Poor
Health problem that limits work	Do you have a serious health problem that limits the amount or kind of work you can do? Y/N	Are you limited in any way in any activities because of physical, mental, or emotional problems? Y/N
Obese	Based on weight/height we compute a mother's BMI	Based on weight/height we compute a mother's BMI
Has health insurance	Are you currently covered by Medicaid or by another public, federal, or state assistance program, or by a private health insurance plan? Y/N	Do you have any kind of health care coverage, including health insurance, prepaid plans such as HMOs, or government plans such as Medicare? Y/N
<u>Health behaviors</u>		
Smokes	In the past month did you smoke cigarettes? Y/N	Smoking status? A. Never smoked B. Former smoker C. Now smokes some days D. Now smokes everyday
Binge drinking	What is the largest number of drinks you had in any single day during the past twelve months? A. 0 B. 1-3 C. 4-10 D. 11-20 E. >20	Do you have four or more drinks on one occasion? Y/N
Drug use	Did you use any of the drugs in list (see footnote 9) "on your own", during the past twelve months? Y/N	
<u>Mental health</u>		
Depressed	Clinically depressed: This assessment is conducted by specialists (see footnote 11)	For how many days during the past 30 days was your mental health (including stress, depression, and problems with emotions) not good? A. 0 B. # of days (from 1 to 30)

Appendix Table 9 - Summary Statistics in FF and BRFSS

Variable	FF	BRFSS¹
Age	26.9 (6.2)	36.4 (8.8)
<u>Race:</u>		
White	0.284	0.404
Black	0.357	0.172
Hispanic	0.290	0.337
Other race	0.070	0.080
Immigrant	0.239	
<u>Education:</u>		
<HS	0.271	0.195
HS	0.329	0.242
Some college	0.203	0.246
College or >	0.197	0.316
<u>Marital Status:</u>		
Married	0.519	0.611
Cohabiting	0.233	0.061
Single²	0.248	0.326
<u>Income^{3,4}:</u>		
1	0.254	0.237
2	0.276	0.192
3	0.136	0.244
4	0.334	0.253
<u>Employment status:</u>		
Employed	0.478	0.570
Unemployed	0.175	0.092
Out of Labor Force	0.344	0.337
<i>N pooled sample</i>	7,080	4,921

Note:

In FF, all characteristics are measured at baseline; in BRFSS these are measured in the pooled sample.

Numbers are weighted using baseline city weights in person specific weights in BRFSS.

¹ Sample includes mothers with children born between 1998 and 2000, in the urban areas of FF baseline states (center city of MSA). Given that child's age (and urban indicator) is only available since 2005, the sample is restricted to 2005-2010 and 13 states (TN and MA not included)

² In BRFSS single includes single, divorced, and widowed women.

³ In FF income levels are: 1) <100% income-to-needs ratio; 2) 100-200%; 3) 200-300%; 4) >=300%

⁴ In BRFSS income levels are: 1) Less than \$20,000; 2) between \$20,000 and \$35,000; 3) between \$35,000 and \$75,000; 4) more than \$75,000.

Appendix Table 10 - The Effect of State UR on Mother's Labor Market Outcomes in FF

	Unemployed	Employed	Out of Labor Force
All	1.056 [0.751]	0.918 [-1.358]	1.071 [0.941]
White	0.995 [-0.02]	1.075 [0.50]	0.832 [-1.13]
Non-Whites	1.066 [0.79]	0.884 [-1.74]	1.117 [1.33]
Married	0.541 [-2.53]	1.105 [0.65]	1.262 [1.32]
Unmarried	1.173 [1.96]	0.989 [-1.50]	0.995 [-0.06]
More than HS	0.884 [-0.75]	0.953 [-0.41]	1.157 [1.02]
HS or less	1.099 [1.12]	0.909 [-1.25]	1.032 [0.37]
N	7,076	7,076	7,076
N changers	704	922	650
Indiv. controls	Y	Y	Y
Bline state FE	Y	Y	Y
Year FE	Y	Y	Y

Note:

Each coefficient comes from a separate regression of Logistic-fixed effects estimates.

T-statistics are shown in brackets; bold font indicates that the result is statistically significant at the 95% level of confidence.

**Appendix Table 11 – The Propensity to Attrit in Year 9 Explained by the UR in Year 5
and Women’s Characteristics**

	(1)	(2)
UR	1.711	5.291
	[2.061]	[1.001]
UR * Mother's Age <19		1.001
		[0.006]
UR * Mother's Age 20-23		0.979
		[-0.208]
UR * Mother's Age 24-27		0.991
		[-0.118]
UR * Mother's Age 28-32		1.001
		[0.021]
UR * Mother is Black		0.821
		[-1.187]
UR * Mother is Hispanic		0.774
		[-1.129]
UR * Mother is other race/ethn		0.787
		[-1.693]
UR * Mother is immigrant		1.062
		[0.338]
UR * Mother is single		0.814
		[-1.015]
UR * Mother cohabitates		0.894
		[-0.616]
UR * Mother's Educ HS		0.905
		[-0.992]
UR * Mother's Educ Some College		0.819
		[-3.733]
UR * Mother's Educ College		0.759
		[-1.131]
UR * Mother's income-to-needs ratio <100%		1.014
		[0.150]
UR * Mother's income: 100-199%		1.099
		[0.718]
UR * Mother's income: 200-299%		1.111
		[0.662]
UR * Child's Age in months		0.988
		[-0.510]
	N	3,797
	3,797	3,797
Indiv. controls	Y	Y
Blime state FE	Y	Y
Year FE	Y	Y

Note:

Each column is a separate regression.

Sample includes year 9 data as a cross section.

The state UR is measured in year 5.

All covariates (mother's age, race/ethnicity, education, marital status, income) are measured at baseline. Errors are clustered at the state level.

T-statistics are shown in brackets; bold font indicates that the result is statistically significant at the 95% level of confidence.

Appendix Table 12 - The Propensity to Attrit in Year 9 Explained by UR and Mother's Health in Year 5

	(1)	(2)
UR	1.727 [1.754]	1.747 [1.765]
UR * Health is Excellent or V. good		0.989 [-0.122]
UR * Health problem limits work		0.749 [-1.605]
UR * Obesity		1.321 [4.201]
UR * Any health insurance		0.882 [-1.920]
UR * Smokes		1.088 [1.036]
UR * 4>= Drinks in 1 Occasion		0.916 [-0.740]
UR * Uses drugs		1.207 [1.777]
UR *Clinically depressed		1.016 [0.114]
<hr/>		
N	3,472	3,472
Indiv. controls	Y	Y
Blinc state FE	Y	Y
Year FE	Y	Y

Note:

Each column is a separate regression.

The state UR and all health variables (health status, health problem that limits work, obesity, etc.) are measured in year 5.

All covariates (mother's age, race/ethnicity, education, marital status, income) are measured at baseline. Errors are clustered at the state level.

T-statistics are shown in brackets; bold font indicates that the result is statistically significant at the 95% level of confidence.